

the elastic modulus also increases to minimize changes in deformation. This suggests that the materials will maintain their shape when subjected to a wide range of externally applied stresses. What's more, the polymers' highly tunable rigidity means that tiny quantities of polymer could be used to make materials that have a wide range of stiffnesses.

Kouwer and colleagues' polymers most closely mimic those found in intermediate filaments (Fig. 1), a class of intracellular polymer that is crucial for cell adhesion and migration and for maintaining cell shape⁴. It will be exciting to see if the authors' approach, or other approaches for making semi-flexible polymers, can be expanded to make synthetic mimics of DNA, actin filaments and microtubules. Another challenge will be to find a way of adding mechanochemically active components⁵ — those that transform chemical energy into mechanical work — to the polymer. This would enable filaments to be made that exhibit exotic polymerization behaviour, such as treadmilling (in which one end of a filament grows while its other end shrinks), or which

NATURE.COM

For more on synthetic

polymers, see:

go.nature.com/xbhhtj

create dynamic instabilities or crosslinks, to form the basis of a molecular motor.

The ability to build 'active' soft materials that

respond to external chemical and mechanical signals will provide opportunities in the areas of condensed-matter physics and materials science for years to come. Such materials might allow the construction of artificial cells and tissues that are more closely compatible physiologically with their counterparts in humans than currently available materials, so that they might be used in the next generation of drug-delivery and tissue-engineering technologies. Active soft materials might also change the way in which we engage with the physical world, by forming the basis of highly responsive and malleable materials and machines. Kouwer and co-workers' polymers are an exciting first step in these directions. ■

Margaret Lise Gardel is at the Gordon Center for Integrated Science, Department of Physics, University of Chicago, Illinois 60637, USA.
e-mail: gardel@uchicago.edu

1. Fletcher, D. A. & Mullins, R. D. *Nature* **463**, 485–492 (2010).
2. Kouwer, P. H. J. *et al.* *Nature* **493**, 651–655 (2013).
3. Gardel, M. L. *et al.* *Meth. Cell Biol.* **89**, 487–519 (2008).
4. Goldman, R. D. *et al.* *J. Struct. Biol.* **177**, 14–23 (2012).
5. Fletcher, D. A. & Geissler, P. L. *Annu. Rev. Phys. Chem.* **60**, 469–486 (2009).

CONDENSED-MATTER PHYSICS

Hidden is more

Physicists have puzzled over a hidden electronic order in a uranium-based material for decades. A new theory attributes it to not just a single but a double breaking of time-reversal symmetry. [SEE ARTICLE P.621](#)

QIMIAO SI

A magnet sticks to a fridge door, but an aluminium spoon does not. This distinction is well understood in terms of the different ways in which the many billions of billions of electrons are collectively organized inside these materials. In a magnet, the electrons form an order: their tiny spins line up along a particular direction, producing an aggregate magnetic moment, whereas in aluminium these spins are randomly oriented. On page 621 of this issue, Chandra *et al.*¹ propose a different kind of electronic order, which could resolve a riddle that has confounded physicists for more than a quarter of a century.

Much of the fascination and challenge of condensed-matter physics lies in figuring out how the electrons are organized in their microscopic world to produce the macroscopic properties observed in the laboratory. The tendency of the electrons in a magnet to develop

order is analogous to that of water molecules to form a rigid spatial pattern as the liquid freezes into ice. This electronic order, called ferromagnetism, breaks time-reversal symmetry: if the time direction were reversed, so would be the direction of the magnetic moment.

The condensed-matter system studied by Chandra and colleagues is URu₂Si₂. This uranium-based compound is a member of a broad class of materials called strongly correlated electron systems, in which a large Coulomb repulsion between the electrons tends to produce spectacular physical phenomena — such as the high-temperature superconductivity observed in copper-based ceramics. This large repulsion in strongly correlated electron systems contrasts with the weak interactions found in many of the materials used in technology, such as silicon, aluminium or even ordinary magnets.

In the mid-1980s, researchers discovered^{2–4} clear signatures of an electronic order in



50 Years Ago

Living with the Atom. By Prof. Ritchie Calder — The author gives his ... contributions to a discussion on responsible reporting. It is difficult for the reporter to steer a course between the attractive liveliness that approaches the sensational, and the dry factual report that few will read ... There is a fear and distrust of scientists, as people who ought to be above human fallibility but unforgivably err like everyone else; and he points out the dangers also of the responsibility for major decisions resting in the hands of men who do not know sufficient about science to be able to challenge with any confidence the advice that comes to them from their experts. There is something in this, although one would feel that a knowledge of men and a flair for consulting the right experts is what brings men to high office.

From *Nature* 2 February 1963

100 Years Ago

'Luminous halos surrounding shadows of heads' — The phenomenon referred to ... can also be seen on grass when the sun is low in the sky ... If the grass surface is near to the observer, a faint halo is seen to surround the shadow of his head, and this is more easily perceived if he is moving than if standing still; my attention was indeed first attracted to this phenomenon when bicycling. J. Evershed

I happened to be watching our shadows as we passed along the edge of a field of young green wheat, when, to my surprise, I noticed a halo of light round the shadow of my own head and neck ... The fact that each observer sees only his own halo obviously precludes this phenomenon from having been the origin of the halos recorded in sacred writings round the head of Christ and others. L. L. Fermor

From *Nature* 30 January 1913