

On the Intelligence of Dogs

WHEN reading in NATURE of November 12, 1885, the abstract of Sir John Lubbock's paper "On the Intelligence of Dogs," I called to mind an incident of a little Blenheim spaniel which belongs to my mother.

The readers of NATURE may perhaps be a little tired of stories relating to the intelligence of the dog, especially when these are illustrations of the effects of training. My excuse for troubling you now is that the following incident seems to indicate a singular power of reasoning.

"Middy" was about nine months old when he was picked off the streets of Melbourne, and he had many traits of the "larrikins," as the human waifs there are called. He had been three months in our family, and we had almost begun to despair of breaking him in to civilised life.

One Sunday my sisters set off for Sunday-school, and were surprised, on nearing the church, to find "Middy" at their heels. He was told to "go home," and he was found at the house on their return. Nothing more was said on the subject, which was forgotten by the next Sunday. But when my sisters entered the school-room on that day, great was their amusement to see the little dog seated calmly as a scholar in one of the classes! He behaved quite quietly during the lessons, and then left with the children, and trotted home alone. To prevent constant repetitions of this behaviour, he had to be caught hours before school-time and shut up. He was very clever in evading capture—crept into hiding early in the day, and bolted when we were off guard. On these occasions he was certain to be found in his place at school.

It perhaps should be especially noted that "Middy" had never been to the church before, and that a whole week had elapsed between his first and second attempts.

MARY KNOTT

7, Kaga Yashiki, Tokio, Japan, January 20

Frost in Devonshire

THE Rev. A. D. Taylor, Rector of Church Stanton, a parish in Devonshire, some 900 feet above sea-level, writes me under date of the 22nd inst. :—

"We have had for three days the most wonderful rime. The trees have been covered, every twig and bud, with ice, on the average an inch at least in depth. I have measured several pieces, and have found them $1\frac{1}{4}$ to $1\frac{3}{4}$ inches from base to edge. The whole place has been like fairy-land, or a silver country. To-day it has all fallen, with a continuous rushing and rattling on the bushes for four hours. The very leaves of the laurels were so frozen that you could take off each leaf a perfect *ice-leaf*—an exact reproduction in transparent ice, of about twice the thickness of this (ordinary letter) paper, of the laurel leaf—every vein and unevenness of edge distinct and clear. The children collected scores of them, and very lovely they looked. I have never seen anything of the sort which would compare with it. The people call it *rängling* (phonetic spelling), a queer word of which I never heard before."

Keen frost in an excessively moist air no doubt sufficiently explains the beautiful phenomenon itself; but can any Devonshire man explain the country people's word?

Bregner, Bournemouth, February 24 HENRY CECIL

"Pictorial Arts of Japan"

IN my review last week of Mr. Anderson's "Pictorial Arts of Japan" I inadvertently wrote the "eight Nirvanas" of Gautama instead of the "eight incidents (more properly 'features'—*fa siang*) of the Nirvana."

F. V. DICKINS
University of London, Burlington Gardens, W., March 1

DISCOVERY OF A NEW ELEMENT BY CLEMENS WINKLER¹

IN the summer of 1885 a rich silver ore was found at Himmelsfürst, near Freiberg; it was pronounced by A. Weisbach to be a new mineral, and was named *Argyrodite*. T. Richter examined its behaviour in the blow-pipe flame, and found that it consisted chiefly of sulphur and silver together with a little mercury, which latter element has never before been found at Freiberg.

¹ From the *Berichte* of the Berlin Chemical Society, No. 3.

The author has analysed the new mineral, and finds that the amount of mercury only amounts to 0.21 per cent., whilst silver is present to the extent of 73-75 per cent., and sulphur to the extent of 17-18 per cent. He also finds a very small quantity of iron, and traces of arsenic. However often and however carefully the analysis was conducted, a loss of 6-7 per cent. always remained unaccounted for. After a long and laborious search for the source of this error, Clemens Winkler has at length succeeded in establishing the presence of a new element in argyrodite. *Germanium* (symbol Ge), as the new element is called, closely resembles antimony in its properties, but can, however, be sharply distinguished from the latter. The presence of arsenic and antimony in the minerals accompanying argyrodite, and the absence of a method of sharply separating these elements from germanium, made the discovery of the new element extremely difficult.

The author, having a more detailed communication in view, confines himself to the following particulars :—

When argyrodite is heated out of contact with the air, which is best effected in a current of hydrogen, a black crystalline and moderately volatile sublimate forms, which melts to brownish-red drops, and which consists principally of germanium sulphide, together with a little mercury sulphide. Germanium sulphide dissolves readily in ammonium sulphide, and, on the addition of hydrochloric acid, is thrown down again in a pure state as a snow-white precipitate, which is immediately dissolved when treated with ammonia; the presence of arsenic or antimony colours the precipitate more or less yellow.

On heating germanium sulphide in a current of air, or on warming it with nitric acid, a white oxide is produced which is not volatile at a red heat and which is soluble in potash solution; when the alkaline solution is acidulated and submitted to the action of sulphuretted hydrogen, the characteristic white precipitate is produced.

The oxide is readily reduced by hydrogen, whilst the sulphide on account of its volatility is more difficult to reduce. The element, like arsenic, has a gray colour and moderate lustre, but is volatile only at a full red heat, and is decidedly less volatile than antimony. Its vapour condenses to small crystals recalling those of sublimed iodine; these show no tendency to melt and could not be confounded with antimony.

When germanium or its sulphide is heated in a current of chlorine it yields a white chloride which is more readily volatile than antimony chloride; its acidulated aqueous solution yields a white precipitate with sulphuretted hydrogen.

The author intends to undertake the determination of the atomic weight of germanium, even if it can be decided only approximately, as this will show whether the new element is to occupy the vacant position in the periodic system between antimony and bismuth.

THE STORY OF BIELA'S COMET¹

II.

BRANDES, one of the two German students spoken of, was riding in an open post-waggon on the night of Dec. 6, 1798, and saw and counted hundreds of these shooting-stars or meteors. At times they came as fast as six or seven a minute. These meteors which Brandes saw that night we know now were bits from Biela's comet. In November 1833 occurred the famous star-shower, which some of you saw. The facts of that shower gave to two New Haven men, Profs. Twining and Olmsted, the clue to the true theory of the shooting-stars. From that date

¹ A Lecture delivered by Prof. H. A. Newton, on March 9, 1874, at the Sheffield Scientific School of Yale College, U.S. From the *American Journal of Science*. Continued from p. 395.