

LETTERS TO THE EDITOR.

The Editor does not hold himself responsible for opinions expressed by his correspondents. Neither can he undertake to return, or to correspond with the writers of, rejected manuscripts intended for this or any other part of NATURE. No notice is taken of anonymous communications.]

Professor Boltzmann's Letter on the Kinetic Theory of Gases.

IN common, I am sure, with all the physical readers of NATURE, I have read Herr Boltzmann's letter with great interest. And I am glad to observe that, though he appears to think I differ from him, that part of his letter which chiefly deals with my criticism on Dr. Watson's idea of what "Boltzmann's Minimum Theorem" is, is simply putting forward, with all his great authority, the view for which I contended. But it is a little hard that Dr. Boltzmann should represent me as endeavouring to *disprove* his theorem when I expressly stated that while I did not know *his* proof, I supposed that it was all right. True, I said that I found it hard to conceive how any proof on the lines of Dr. Watson's could be valid because that proof appeared to me to be a purely dynamical proof, and I applied the reversibility argument to show that a purely dynamical proof was impossible, so that the H-theorem could not be a purely dynamical theorem; and after indicating the lines on which it appeared that there might be an average dynamical theorem, I asked if some one would say what the H-theorem really was.

Thereupon Mr. Burbury wrote a helpful letter, which he followed up by a still more helpful correspondence, in which verbal misunderstandings were gradually cleared away, which showed that the proof of the H-theorem considered as a dynamical theorem, not as a theorem in probabilities, assumed that in one respect the configuration was, before each set of collisions, already perfectly average, and that this condition is violated in the reversed motion; so that the theorem, regarded as a dynamical theorem, is not proved for configurations in general, but for those possessing a certain amount of "average" already—a restriction which comes to the same thing as the limitation imposed by Prof. Boltzmann when he says *the theorem is not a dynamical theorem, but one in probabilities*.

Shortly after Mr. Burbury's letter appeared, Dr. Watson wrote denying that the criticism from reversibility applied, and claiming that the theorem was a general dynamical theorem, in the sense that it applied to *all* configurations. Enlightened by Mr. Burbury, I now see that Dr. Watson's reasoning is not open to the objection that it proves a general dynamical theorem; but I cannot blame myself for thinking that it did, for that was what Dr. Watson himself believed it to do, and what his language naturally implies. Moreover, after perceiving the oversight which vitiates the proof in its present form, I did not examine it further.

Prof. Boltzmann has misunderstood Mr. Burbury and me in one or two particulars. He denies that there are as many configurations for which dH/dt is positive as there are for which it is negative. He evidently thinks that we mean something different from the bare meaning of the words, which are certainly true. It is easy to explain what we do not mean (I say *we*, for I am sure Mr. Burbury will agree with me). Suppose $H=10$ to be the minimum value of H for a given system of molecules, we do not mean that among all the configurations for which $H=50$, there are as many which will, if left to themselves, turn into configurations for which $H=60$, as will turn into configurations for which $H=40$. The illustration, which to my mind has most clearly removed the apparent contradiction in the statement that there are as many configurations for which H will increase as decrease, while yet the probability is that H will on the whole decrease, is that of a y turned upside down, thus Λ . For every downward path there is an upward path, *i.e.* the reversed direction; yet starting from the angle there are two ways down for one way up, so that there is a greater probability of going down than up. If in the reversibility argument one could assert, not merely that there are as many configurations for which H tends to increase as to decrease, but that for any given value of H there were as many configurations which tend to increase as to decrease, then the conclusion that H was as likely to increase as to decrease could be deduced. But the argument is quite invalid when we set off a configuration for which H increases against one for which it decreases, *although*

the values of H for each are different. As an illustration more closely allied to the case of a gas, we might take a tree turned upside down, with an infinite number of branches passing through each point of its substance in all directions, there being at every point more branches tending downward than upward (because those whose tangents are horizontal may be said to tend downward on each side), and every upward branch finally turns downward and tends to become nearly horizontal at last, when H is near its minimum value.

To my mind this appears a far better way of meeting the difficulty than Prof. Boltzmann's illustration of the dice, for so far as I can see, all that he has shown is that if you start from an exceptionally high ordinate, *i.e.* one over the average, you are likely, after a considerable time, to get to lower ordinates in whichever direction you go, and an opponent might answer that if you start from an exceptionally low ordinate you are likely to get higher ones in whichever direction you go, and that there must of course be as many deviations below the average as above it, so that if you start from an arbitrary point in an arbitrary direction, you are just as likely to get to higher as to lower ordinates. In point of fact this appears to be the case for his curve, while it is not true for the tree or for a gas.

Prof. Boltzmann must have put an entirely wrong construction on something or other, which I suppose I have written, when he says I object to the Maxwell Law of distribution because it would ultimately lead to the total kinetic energy of the universe being equally distributed among every degree of freedom of every particle in the universe. Instead of considering that to be *à priori* improbable, I hold exactly the view put forward by Prof. Boltzmann.

With regard to the first portion of Prof. Boltzmann's letter, there is so much that is speculative in it that any discussion would occupy more space than I feel entitled to claim. I will only say that the idea that a gas takes years to come to thermal equilibrium seems hardly consistent with vibrational portion of the kinetic theory being of practical value, when applied to gas which has only had a few hours to settle down.

EDWARD P. CULVERWELL.

Trinity College, Dublin, March 6.

IT seems to me that my meaning has not been expressed quite clearly; therefore, it may be worth while to add one remark. Not for every curve, but only for the particular form of the H-curve, disymmetrical in the upward and downward direction, can it be proved that H has a tendency to decrease. This particular form is very well illustrated by Mr. Culverwell's suggestion of an inverted tree. The H-curve is composed of a succession of such trees. Almost all these trees are extremely low, and have branches very nearly horizontal. Here H has nearly the minimum value. Only very few trees are higher, and have branches inclined to the axis of abscissæ, and the improbability of such a tree increases enormously with its height. The difficulty consists only in imagining all these branches infinitely short.

Finally there is the difference between the ordinary cases, where H decreases or is near to its minimum value, and the very rare cases, where H is far from the minimum value and still increasing. In the last cases, H will reach, probably in a very short time, a maximum value. Then it will decrease from that value to the well-known minimum value.

Paris, April 6. LUDWIG BOLTZMANN.

The Recent Auroral Phenomenon.

ON the evening of March 13, from 7.35 to 8.5, Greenwich mean time, I was a spectator of the abnormal display of Aurora Borealis which attracted so much attention at various places throughout the country. It appeared here as a belt of light spanning nearly the whole sky in a great circle from east to west. When first noticed by me at 7.35, the streak extended from near the hind quarters of Leo to the head of Aries, or from R.A. 169°, Decl. + 16° to R.A. 24°, Decl. + 22°.

At the time the streak was altogether cometary in appearance, beginning in a fine point, but it gradually changed in form, moving at the same time towards the south. Eventually it also shortened so considerably that just before my last view of it, it only extended from γ Geminorum to γ Ceti. Its greatest breadth was about 12°.