

is thus doing a service to the nation, as well as extending interest in natural knowledge. Dr. P. Bedson, professor of chemistry at the Durham College of Science, has, we are glad to see, recently shown the Economic Society of Newcastle-on-Tyne some of the lessons taught by the growth of science and industry in Germany during the present century. A reasonable and organised system of education, and schools in which students receive a thorough grounding in the principles of science, and afterwards contribute something to the advancement of knowledge, are the chief factors in Germany's industrial progress. Referring to the system of examinations which still dominates so much of our educational work, and finds its highest development in connection with university teaching, Prof. Bedson points out that it partakes of the character of the training of a stud of racers. He adds:—"Possibly the instinct of sport, so characteristic of the English people, it is which commends the system of competitive examination. Too much is made of what should be regarded as a minor duty of the University, viz. the testing and marking of its students, and too little of the higher function, the training of students under first-rate teaching, with the object that those so trained should help forward the advancement of knowledge." It is satisfactory to know that the movement in favour of rational teaching in elementary schools, and regard for research in institutions of university rank, is gradually affecting scientific education in this country.

SCIENTIFIC SERIALS.

Symons's Monthly Meteorological Magazine, July.—This number contains the completion of two interesting papers, by Mr. E. D. Archibald, on Indian famine-causing droughts, and their prevision. The principal facts are summarised as follows: (1) Extensive droughts occur in the dry area of Southern India at intervals of nine to twelve years, and usually, but not regularly, about a year before the sun-spot minimum. When the conditions are sufficiently acute, famine occurs in the following year. (2) A severe drought in the peninsular of Southern India is followed by a severe drought and ensuing famine in Northern India in about five cases out of seven. (3) Summer droughts tend to occur in Northern India in years of maximum sun-spot, connected in some way with the abnormal high pressure over Western Asia which prevails at such epochs. There is thus a double periodicity of droughts and famine in North India, and a single periodicity in South India in the sun-spot cycle, though the relation between the phenomena is too spasmodic and irregular to be utilised as a trustworthy factor for prevision.

Annalen der Physik, No. 6.—Interruption spark in the alternating circuit with metallic electrodes, by L. Kallir. The author shows that the impossibility of producing an alternate-current arc between metallic electrodes is due to the fact that the spark is confined to one semi-period of the current. Or if it extends over several periods, it is intermittent, and only appears at every alternate semi-period.—Thermoelectric force of some metallic oxides and sulphides, by A. Abt. Pyrolusite, pyrrhotite, pyrites, and chalcopyrite were used in conjunction with various metals or with each other. A pyrites-chalcopyrite couple gave an E.M.F. 10·8 times as high as an antimony-bismuth couple under the same conditions.—Anomalous electromagnetic rotatory dispersion, by A. Schmauss. Measurements of the Faraday effect for various wave-lengths in fuchsine solutions and in didymium glass justify the general conclusion that optical anomaly in dispersion is invariably associated with a corresponding electromagnetic anomaly. In strongly absorbing media the anomaly extends for a considerable distance on both sides of the absorption band, and it increases with the concentration and with the narrowness and sharpness of the absorption band.—Point discharges, by E. Warburg. In carefully purified nitrogen, the current intensity obtained from the discharge of a fine point charged to - 3310 volts is 200 times as great as from a point charged to + 5180 volts. A slight admixture of oxygen reduces the proportion to 4:1.—Band spectrum of aluminium, by G. A. Hemsalech. The author quotes some experiments which go to show that the band spectrum of aluminium is due, not to the oxide, but to the metal itself.—Behaviour of radium at low temperatures, by O. Behrendsen. Cooling a radium preparation down to the temperature of liquid air reduces its activity by 50 per cent. Restoration to the ordinary temperature produces a considerable but transient increase of activity.—Production of kathode rays by ultra-violet light, by P. Lenard. The discharge of electrified

bodies by ultra-violet light is due to their emitting kathode rays when the ultra-violet light impinges upon them. The author exhausted a vacuum tube until it no longer allowed any discharge to pass. He then exposed the kathode to ultra-violet rays from a zinc spark gap. The discharge set in again immediately, but no discharge could be obtained by similarly illuminating the anode. The rays which produce the discharge across the absolute vacuum can be deflected by a magnet, and their velocity is about one-thirtieth of the velocity of light.

SOCIETIES AND ACADEMIES.

LONDON.

Royal Society, February 8.—"On Electric Touch and the Molecular Changes produced in Matter by Electric Waves." By Jagadis Chunder Bose, M.A., D.Sc., Professor of Physical Science, Presidency College, Calcutta. Communicated by Lord Rayleigh, F.R.S.

It is claimed that the experiments described in the paper show:—

- (1) That ether waves produce molecular changes in matter.
- (2) That the molecular or allotropic changes are attended with changes of electric conductivity, and this explains the action of the so-called coherers.
- (3) That there are two classes of substances, positive and negative, which exhibit opposite variations of conductivity under the action of radiation.
- (4) That the production of a particular allotropic modification depends on the intensity and duration of incident electric radiation.
- (5) That the continuous action of radiation produces oscillatory changes in the molecular structure.
- (6) That these periodic changes are evidenced by the corresponding electric reversals.
- (7) That the "fatigue" is due to the presence of the "radiation product," or strained B variety.
- (8) That by means of mechanical disturbance or heat, the strained product can be transformed into the normal form, and the sensitiveness may thereby be restored.

June 21.—"An Experimental Investigation into the Flow of Marble." By Frank D. Adams, M.Sc., Ph.D., Professor of Geology in McGill University, Montreal, and John T. Nicolson, D.Sc., M.Inst.C.E., Head of the Engineering Department, Municipal Technical School, Manchester. Communicated by Prof. H. L. Callendar, F.R.S.

The following is a summary of the results arrived at:—

- (1) By submitting limestone or marble to differential pressures exceeding the elastic limit of the rock and under the conditions described in this paper, permanent deformation can be produced.
- (2) This deformation, when carried out at ordinary temperatures, is due in part to a cataclastic structure and in part to twinning and gliding movements in the individual crystals comprising the rock.
- (3) Both of these structures are seen in contorted limestones and marbles in nature.
- (4) When the deformation is carried out at 300° C. or better at 400° C., the cataclastic structure is not developed, and the whole movement is due to changes in the shape of the component calcite crystals by twinning and gliding.
- (5) This latter movement is identical with that produced in metals by squeezing or hammering, a movement which in metals, as a general rule, as in marble, is facilitated by increase of temperature.
- (6) There is therefore a flow of marble just as there is a flow of metals, under suitable conditions of pressure.
- (7) The movement is also identical with that seen in glacial ice, although in the latter case the movement may not be entirely of this character.
- (8) In these experiments the presence of water was not observed to exert any influence.
- (9) It is believed, from the results of other experiments now being carried out but not yet completed, that similar movements can, to a certain extent at least, be induced in granite and other harder crystalline rocks.

"On the Effects of Changes of Temperature on the Elasticities and Internal Viscosity of Metal Wires." By Andrew Gray, LL.D., F.R.S., Professor of Natural Philosophy in the University of Glasgow, and Vincent J. Blyth, M.A., and