

Calendar of Industrial Pioneers.

October 1, 1838. Charles Tennant died.—The founder in 1797 of famous chemical works at St. Rollox, Glasgow, Tennant while manager of a bleaching field near Paisley discovered a method of controlling chlorine gas by the admixture of lime. He introduced the manufacture of chloride of lime in a solid state, to which he gave the name bleaching powder. His production of bleaching powder in 1799–1800 was 52 tons, the price being 140*l.* per ton. By 1835 the St. Rollox works had become the most important chemical works in the world.

October 2, 1804. Nicolas Joseph Cugnot died.—A military engineer and the author in 1766 of "Éléments de l'art militaire ancien et moderne," Cugnot in 1769 made the first steam-propelled road carriage, and two years later built a steam tractor for the French Government for hauling artillery. This vehicle was to carry a load of 4½ tons at 2¼ miles per hour. Though never used, this carriage is preserved in the Conservatoire des Arts et Métiers.

October 3, 1867. Elias Howe died.—One of the chief pioneers of the sewing machine, Howe was the son of a farmer of Spencer, Massachusetts, and was born in 1819. He began work on the sewing machine in 1841, took out a patent in 1846, and was one of the first inventors to place the eye of the needle towards the point.

October 4, 1821. John Rennie died.—Acknowledged as the greatest civil engineer of his day, Rennie was the builder of the London Docks, the East India Docks, the Plymouth Breakwater, Waterloo and Southwark Bridges, and he prepared designs for London Bridge. He was born at Phantassie, East Lothian, in 1761, gained practical experience under Andrew Meikle, attended the lectures of Robison and Black, and in 1789 erected the Albion Mills for Boulton & Watt, in London, the site of which was afterwards occupied by Rennie's workshops. He is buried in St. Paul's Cathedral.

October 5, 1892. Alexander Carnegie Kirk died.—The author of many improvements in marine engineering, Kirk, after gaining experience at Maudslay's and at Elder's, became a partner in 1877 in the firm of Napier. He was especially known for his advocacy of high steam pressure and the triple-expansion engine, the advantages of which were demonstrated in the s.s. *Aberdeen* built by him in 1882, which on a voyage to Australia showed a saving of 500 tons of coal.

October 6, 1905. Charles Brown died.—Brown has been called the founder of mechanical industry in Switzerland. Brought up in London, in 1851 at the age of 24 he entered the service of Sulzer Brothers, a firm of mechanical engineers at Winterthur. He established afterwards the Swiss locomotive works at Winterthur and also played a prominent part in the creation of the Swiss electrical industry.

October 7, 1908. Jean Baptiste Gustave Adolphe Canet died.—A distinguished armament engineer, Canet was trained at the École Centrale des Arts et Manufactures, fought in the Franco-German War, and for a time engaged in railway engineering. From 1872 to 1881 he was associated with Vavasseur at the London Ordnance Works, and in 1876 brought forward his theory of hydraulic brakes for checking the recoil of guns. Returning to France he became the head of armament works at Havre and after the amalgamation of these works with those of Schneider at Creusot became manager.

Societies and Academies.

SWANSEA.

Institute of Metals, September 20.—G. D. Bengough and J. M. Stuart: The nature of corrosive action, and the function of colloids in corrosion (Sixth Report to the Corrosion Research Committee of the Institute).—Sir Henry Fowler: The effect of superheated steam on non-ferrous metals used in locomotives. Superheated steam as used on locomotives generally leaves the superheater at a temperature of 340° C. On the Midland Railway, piston tail rod bushes were made of M.R. A.1 alloy (copper, 87; tin, 9; zinc, 2; lead, 2). A phosphor bronze (copper, 88; tin, 11; phosphorus, 1) has been found satisfactory. For piston rod packing, McNamee rings (copper, 75.5; tin, 8.5; zinc, 0.33; phosphorus, trace; nickel, 0.5; lead, 15.0) are used satisfactorily. These rings prevent the steam coming into contact with the white metal (lead, 70; antimony, 30) packing rings. With the temperature rising to 425° C. the packing rings may fuse. Piston valve fittings and cylinder relief valves are made of alloy M.R. A.1. For by-pass valves which are subjected to shock, a nickel-brass gave good service, but was replaced for economy by malleable iron or steel castings.—A. H. Munday, C. C. Bissett, and J. Cartland: White metals. The manufacture and use of white metal for industrial purposes is described, and constitution and micro-structure are dealt with only so far as the uses and manufacture are concerned. Antifriction or bearing metal, printing alloys, die-casting alloys, metals for chemical works castings, solders, are discussed.—J. H. Andrew and R. Higgins: Grain-size and diffusion. Diffusion at high temperatures may take place simultaneously with grain-growth, while at low temperatures it promotes a breakdown in the grain-size. These results have been applied to the annealing treatment of castings. It has been assumed that in the interior of the crystalline grains the system of closest packing holds, while at the boundaries the atoms in the separate grains touch only at one part of the circumference. This explains the decrease in specific gravity with an increase in the number of grains, for in such an arrangement free spaces occur. Plastic deformation, by shifting some of the atoms from their positions of equilibrium, will cause them to rearrange themselves when heated to a sufficiently high temperature. This rearrangement will be such that the stressed atoms will fall in, row for row, with the unstrained atoms of the adjacent crystal. This effects a gradual migration of the grain boundary which, proceeding from every side of a crystalline unit, may result in one grain being divided up and absorbed by others. The final bounding surface will result when the boundary configuration is reached.

PARIS.

Academy of Sciences, August 28.—M. L. Maquenne in the chair.—L. Mangin and N. Patouillard: The destruction of the woodwork at the château of Versailles by *Phellinus cryptarum*. A detailed examination of the oak beams showed a varied fauna and flora, but *Phellinus cryptarum* is mainly responsible for the damage. This fungus has not hitherto been regarded as destructive to wood.—Jacques

Chokhate: The development of the integral $\int_a^b \frac{p(v)}{x-y} \cdot dy$ as a continued fraction.—Ch. N. Moore: The equivalence of the methods of summation of Cesàro and of Holder for multiple limits.—Nilos Sakellariou: Polar systems.—Amédée Béjot: Placing in reciprocal

E. C. S.

© 1922 Nature Publishing Group

perspective, figures of the same species.—M. Gignoux and P. Fallot: The marine quaternary on the Mediterranean coasts of Spain.—Raoul Combes and Denise Kohler: The rôle of respiration in the diminution of the carbohydrates in leaves during the autumnal yellowing. It has been commonly held that during the change of colour of leaves in the autumn, the carbohydrates are withdrawn from the leaf and stored in the plant as reserves. It has been proved by Michel Durand that some of the carbohydrates are removed by rain, and in the present communication proof is given that part is used up by respiration and leaves as carbon dioxide.—L. Carrere: The sphincter of the iris in the selacians. This muscle in the selacians, especially in species possessing a pupil-shaped opening, is more developed at the nasal and temporal extremities of the pupil: it is less important, and may even disappear, in the ventral and dorsal sectors.—Paul Wintrebert: The mechanical polarity of the germ of selacians (*Scylliorhinus canicula*) at the time of gastrulation.

September 4.—M. L. Guignard in the chair.—Théodore Varopoulos: A theorem of M. Rémoundos.—Alf. Guldberg: The theorem of M. Tchebycheff.—Victor Henri and Pierre Steiner: Absorption of the ultraviolet rays by naphthalene. From a quantitative study of the absorption of solutions in hexane, ether, alcohol, and water, seventeen bands have been found between wave lengths 3207Å and 2563Å. These results are compared with those previously obtained for benzene by a similar method.—Erik Hulthén and Ernst Bengtsson: Researches on the band spectra of cadmium.—G. Murgoci: The classification of the blue amphiboles and of certain hornblendes.—Marcel Mirande: The formation of anthocyanin under the influence of light on the scales of the bulbs of certain lilies.—Raphaël Dubois: The destruction of mosquitos by eels. Goldfish have been suggested for destruction of mosquitos as they eat the larva, but they have the disadvantages of being costly and requiring a pure and well-aerated water. Young eels in the spring are equally voracious and devour the larvæ readily. They are more readily procurable than goldfish, and live equally well in fresh and salt water, and even in water containing sewage effluent.

SYDNEY.

Royal Society of New South Wales, August 2.—Mr. C. A. Sussmilch, president, in the chair.—C. E. Fawsitt and C. H. Fischer: The miscibility test for eucalyptus oils. Instead of testing the solubility by measuring the volume of aqueous alcohol required to obtain complete solution of a measured volume of oil, the critical solution temperatures with definite mixtures of alcohol and water are taken after the manner of testing fixed oils. This method is more sensitive for the indication of small changes in composition of the oil. The critical solution temperature in some cases varies markedly with time and as the oil is kept.—R. T. Baker and H. G. Smith: The Melaleucas and their essential oils, Pt. VI. Two species are discussed, *Melaleuca ericifolia*, Sm., and *M. Deanei*, Fr. M. Oil was first distilled from *M. ericifolia* by Mr. J. Bosisto in 1862, and Dr. J. H. Gladstone in 1864 determined its physical constants. The chief oxygenated constituent was thought to correspond with that in ordinary oil of "cajuput." The yield of oil obtained by the present authors was 0.8 per cent., and the chief oxygenated constituent found to be dextrorotatory terpineol, while less than 10 per cent. of cineol was present. Pinene, limonene, and a sesquiterpene were also detected. The yield of oil from young material of *M. Deanei* was also 0.8 per cent., and consisted almost entirely of pinene

with about 15 per cent. of cineol. Old leaves of this species contain very little oil.—A. R. Penfold: The essential oil from *Bachhousia myrtifolia*, Pt. I. This small tree inhabits gullies containing running water in the coast and coast mountain districts of New South Wales. Material collected at Lane Cove near Sydney, and at Currowan of the southern district, yielded 0.3-0.75 per cent. of a brown-yellow oil, varying with the time of year and locality. The oil possesses a pleasant odour and is heavier than water. Its principal constituent is elemicin (80 per cent.), a somewhat rarely occurring phenol ether. The remainder of the oil consists of α -pinene, unidentified alcoholic bodies and phenols, sesquiterpene, and a paraffin of melting-point 62°-63° C.

CAPE TOWN.

Royal Society of South Africa, August 16.—Dr. J. D. F. Gilchrist, president, in the chair.—W. A. Jolly: The rhythm of discharge of the spinal centres in the frog. The rate of discharge of the cord in Xenopus at different temperatures, as indicated by galvanometric records from the gastrocnemius muscle reflexly excited, was discussed.—J. P. Dalton: On the mathematics of the homogeneous balanced action. It has been shown by the author that the integrated velocity equations of chemical reactions can be written in terms of a certain function. The same function may be employed in the treatment of the homogeneous balanced action.

Official Publications Received.

Western Australia. Annual Progress Report of the Geological Survey for the Year 1921. Pp. 61. (Perth.)
Northampton Polytechnic Institute, St. John Street, London, E.C. Announcements, Educational and Social, for the Session 1922-1923. Pp. 248. (London: Northampton Polytechnic Institute.)
New Zealand. Department of Mines: Geological Survey Branch. Palæontological Bulletin No. 9: The Upper Cretaceous Gastropods of New Zealand. By Dr. Otto Wilckens. Translated into English by the Author. Pp. iv+42+5 plates. (Wellington, N.Z.)
Prospectus of the Royal College of Art, S. Kensington, London. Session 1922-1923. Pp. iv+29. (London: H.M. Stationery Office.) 9d. net.
London School of Tropical Medicine, Department of Helminthology. Collected Papers, 1922 (Part 2), Nos. 16-25. Pp. 3+4+11+7+9+75+3+7+11+5. (London: 23 Endsleigh Gardens.)

Diary of Societies.

MONDAY, OCTOBER 2.

SOCIETY OF ENGINEERS (at Geological Society), at 5.30.—C. H. J. Clayton: The Economics of Arterial Land Drainage.

THURSDAY, OCTOBER 5.

ROYAL AERONAUTICAL SOCIETY (at Royal United Service Institution), at 5.30.—Prof. L. Baird: The Work of S. P. Langley.
CHILD-STUDY SOCIETY (at Royal Sanitary Institute), at 6.—Dr. C. W. Kimmins: Visual Humour.
ROYAL PHOTOGRAPHIC SOCIETY, at 8.—R. H. Lawton: The Use and Misuse of Short Focus Lenses.
CHEMICAL SOCIETY, at 8.—H. Bassett and R. G. Durrant: Cupric Tetrammine Nitrite and the Corrosion of Copper by Aqueous Solutions of Ammonia and of Ammonium Nitrate.—C. K. Ingold and H. A. Piggott: The Additive Formation of Four Membered Rings. Pt. I. The Synthesis and Resolution of some Derivatives of Tetrahydro-1:3-Diazine.

FRIDAY, OCTOBER 6.

JUNIOR INSTITUTION OF ENGINEERS, at 7.30.—W. A. Tookey: Engineering in Bacon Factories.
ROYAL PHOTOGRAPHIC SOCIETY, at 8.—F. Lambert: The Beginnings of London.

PUBLIC LECTURE.

THURSDAY, OCTOBER 5.

KING'S COLLEGE, at 5.30.—Prof. H. Wildon Carr: The New Method of Descartes and the Problems to which it gave rise (I). Succeeding Lectures on Oct. 6, 12, 13, 19, 20, 26, 27, Nov. 2, 3.