

INTRACELLULAR DISTRIBUTION OF RHODANESE

Fraction	Liver (Ludewig and Chanu- tin)	Cardiac muscle (Moyle)		
	Total activity (per cent)	Total activity (per cent)	Total dry wt. (per cent)	Specific activity
Homogenate	100	100	100	1
(a) Residue	15	44		
(b) Large particles	62	2	9.3	0.2
(c) Supernatant	7	56		
a + b + c	84	102		

nuclear residue, (b) sarcosomes and (c) sarcoplasm. Rhodanase activity was estimated by the method of Cosby and Sumner⁶. A comparison of the activity of the homogenate and the tissue fractions showed that the sarcosomes contained no rhodanase, the total activity of the muscle being only about 10 per cent that of liver (per unit weight).

Cleland and Slater² have reported a parallel situation in the distribution of myokinase, which occurs in the sarcoplasm of cardiac muscle but in the mitochondria of liver⁷.

So, although the 'large granules' from liver and heart muscle show similarity in being the site of respiratory enzyme systems, they are not completely identical units differing only slightly in morphology from tissue to tissue. In their diverse non-respiratory functions they may take very different roles in the metabolism of their respective tissues.

I wish to thank Dr. M. Dixon for valuable advice, and the Medical Research Council for a grant.

JENNIFER M. MOYLE

Department of Biochemistry,
University of Cambridge. April 22.

¹ Green, D. E., "Enzymes and Enzyme Systems" (Harvard, 1951).
Hogeboom, G. H., *Fed. Proc.*, **10**, 640 (1951).
Lehninger, A. L., "Enzymes and Enzyme Systems" (Harvard, 1951).

² Cleland, K. W., and Slater, E. C., *Biochem. J.*, **53**, 547 (1953).

³ Paul, M. H., Fuld, M., and Sperling, E., *Proc. Soc. Exp. Biol. Med.*, **79**, 349 (1952).

⁴ Ludewig, S., and Chanutin, A., *Arch. Biochem.*, **29**, 441 (1950).

⁵ Hogeboom, G. H., Schneider, W. C., and Pallade, G. E., *J. Biol. Chem.*, **172**, 619 (1948).

⁶ Cosby, E. L., and Sumner, J. B., *Arch. Biochem.*, **7**, 457 (1945).

⁷ Kielley, W. W., and Kielley, R. K., *J. Biol. Chem.*, **191**, 485 (1951).

Activation of Fumaric Hydrogenase by Ferrous Ions

FISCHER *et al.*¹ described an enzyme from yeast which catalyses the reduction of fumarate to succinate by leuco-dyestuffs. They showed that the coenzyme is flavine adenine dinucleotide.

It has now been found that addition of flavine adenine dinucleotide does not fully restore the activity of this enzyme after dialysis and adsorption on alumina C_γ, but that addition of ferrous ions is also necessary.

Thunberg tubes contained: 0.5 ml. 0.05 M phosphate buffer pH 7.4 and 1 ml. 0.03 M fumarate. In stopper: 0.5 ml. 0.0002 M janus green reduced with 0.2 mgm. hydrosulphite. Gas, nitrogen.

ACTIVATION OF FUMARIC HYDROGENASE

Additions	Oxidation time (min.)
Nil	> 25
Fe ⁺⁺	> 25
Enzyme	10½
Enzyme + Fe ⁺⁺	1½

Additions: 1 ml. enzyme (dialysed in the presence of flavine adenine dinucleotide); 0.2 ml. 0.01 M FeSO₄. Other metals such as Mn⁺⁺ are ineffective. Keilin-Hartree preparations of succinic dehydrogenase from heart muscle also lose activity on dialysis; addition of ferrous ions enables the reduction of fumarate to take place. It would appear, therefore, that ferrous ions are required for electron transfer in succinate-fumarate systems.

K. HARRISON

Department of Biochemistry,
Cambridge. May 4.

¹ Fischer, F. G., *et al.*, *Ann. Chem.*, **552**, 203 (1942).

Chromaffin Bodies of Various Species of Dogfish

In lower vertebrates, structures homologous to the adrenal gland of higher vertebrates remain separated throughout life. In the dogfish, for example, an unpaired inter-renal body representing the adrenal cortex is quite separate from the chromaffin bodies or rudimentary adrenal medulla. Extracts of each component can therefore be made without contamination by the other part.

Since workers in both our departments are interested in the biosynthesis of adrenaline, we have investigated by chromatographic and biological methods the amines present in the chromaffin bodies of *Scylliorhinus canicula* and *stellaris* (in Bari), *Squalus acanthias* and *Mustelis canis* (in Dundee), and *Torpedo marmorata* (in both Bari and Dundee). Our pooled results are shown in the accompanying table. Large quantities of adrenaline and noradrenaline are contained in these extracts; but hydroxytyramine, dihydroxyphenylalanine, dihydroxyphenylserine, tyramine and octopamine (*p*-norsynephrine) are not present in detectable amounts. Concentrated extracts of the livers, kidneys, intestinal tract, and inter-renal bodies of these dogfish species do not contain significant amounts of any of the seven substances mentioned.

AMINES (μGM./GM.) PRESENT IN THE CHROMAFFIN BODIES OF DOGFISH SPECIES

Species	Adrenaline	Noradrenaline	Noradrenaline (per cent in total)
<i>Squalus acanthias</i>	900	2,400	73
<i>Mustelis canis</i>	1,000	2,200	69
<i>Scylliorhinus canicula</i>	1,150	2,200	66
<i>Scylliorhinus stellaris</i>	850	1,900	67
<i>Torpedo marmorata</i>	150	600	80

These results do not assist in advancing the method by which adrenaline and noradrenaline are formed in the body. However, it is evident that even the pathways through dihydroxyphenylalanine and hydroxytyramine, or through tyramine and *p*-norsynephrine, must be considered doubtful.

D. M. SHEPHERD

G. B. WEST

Department of Pharmacology and Therapeutics,
University of St. Andrews Medical School,
Dundee.

V. ERSFAMER

Department of Pharmacology,
University of Bari,
Italy. April 29.