

Extreme weather events

SIR — The seasonally recurring oceanic El Niño phenomenon is associated with extreme weather conditions¹. Positive phases of the tropical sea-surface temperature oscillation have been connected with unusually rainy weather over the southwestern United States, while the negative phases, usually referred to as La Niña, have been plausibly associated with extreme dry conditions over the central United States.

Since the oceanic oscillation was first connected by dynamic and thermodynamic considerations² to the atmospheric Southern Oscillation, understanding of the coupled ocean-atmosphere El Niño/Southern Oscillation (ENSO) phenomenon has increased dramatically¹, leading to hopes that these events could be predicted a year or so in advance. Statistical prediction models of ENSO³ — based on the analysis of sea-surface temperature and surface wind fields — and dynamical models⁴ can forecast events with some confidence out to 6 months or even 1 year.

These models try to forecast the detailed geographical distribution of oceanic and atmospheric variables affected by the ENSO cycle. This is a more difficult task than telling whether an El Niño or La Niña event will occur; univariate indicators of ENSO can be helpful for the latter. The Southern Oscillation index computed from time series of sea-level pressure at Tahiti and Darwin, Australia, has good diagnostic value. Minima in this index are associated with El Niño events, whereas maxima correspond to La Niña events¹.

A combination of singular-spectrum analysis⁵ and the maximum entropy method⁶ seems to hold promise for predicting the ENSO cycle 2–3 years in advance. Singular-spectrum analysis is a variant of principal component analysis applied in the time domain; it filters out variability unrelated to ENSO, separating the quasi-biennial 2–3-year variability from a lower-frequency 4–6-year El Niño–La Niña cycle; the total variance associated with ENSO combines the quasi-biennial and lower-frequency

modes⁷. The corresponding time series is shown in the figure, where it is compared with a 5-month running mean of the original Southern Oscillation index. The additional smoothness of the filtered index reflects that of the underlying temporal principal components and is crucial to the success of the forecasts using the maximum entropy method.

Measured objectively by its correlation with the observed time series in a hindcasting mode, the predictability of the principal components which carry the lower-frequency mode is between 2.5 and 3 years, a little less than the predictability of the more regular quasi-biennial mode. The figure contains a forecast based on data up to the end of February 1992. The Southern Oscillation index is expected to reach its next maximum in early 1994. The drought that affected the continental United States in the summer of 1988 has been related to the 1988–89 La Niña event and to the Southern Oscillation index maximum of January 1989⁸. The next La Niña event, if correctly predicted, could be associated with a drought over the continental United States during the second half of 1993. In addition, an El Niño event is also predicted for 1996–97, when the Southern Oscillation index, as filtered by singular-spectrum analysis, is expected to reach its next minimum.


Christian L. Keppenne*†

Michael Ghil*

*Department of Atmospheric Sciences,
University of California,
Los Angeles, California 90024, USA

†Jet Propulsion Laboratory,
California Institute of Technology,
Pasadena, California 91109, USA

1. Philander, S. G. *El Niño, La Niña, and the Southern Oscillation* (Academic, San Diego, California, 1990).
2. Bjerknes, J. *Int. Trop. Tuna. Comm. Bull.* **5**, 219–271 (1966).
3. Barnett, T. *et al. Science* **241**, 192–196 (1988).
4. Cane, M., Zebiak, S. & Dolan, S. *Nature* **231**, 827–832 (1986).
5. Vautard, R. & Ghil, M. *Physica D35*, 395–424 (1989).
6. Penland, C., Ghil, M. & Weickmann K. M. *J. geophys. Res.* **96**, 22659–22671 (1991).
7. Rasmusson, E. M., Wang, X. & Ropelewski, C. F. *J. mar. Syst.* **1**, 71–96 (1990).
8. Mo, K. C., Zimmerman, J. R., Kalnay, E. & Kanamitsu, M. *Mon. Weath. Rev.* **119**, 1512–1532 (1991).



Southern Oscillation index series obtained by singular-spectrum analysis (solid, thin line); 5-month running mean of the index (dotted, thin line); and forecast obtained by applying the maximum-entropy method to the leading temporal principal components of the index (solid, thick line).

Cosmic microwave background

SIR — Fluctuations of the cosmic microwave background from the COBE satellite (discussed recently in *News and Views*¹) have been widely interpreted as providing confirmatory evidence for the Big Bang explanation of this radiation, the fluctuations being taken to represent the signatures of galaxy formation in the early Universe. Alternative explanations for the microwave background, involving thermalization, can also account for the observed energy density of the radiation and can also produce intensity fluctuations of the kind that were observed by COBE.

For such a model involving thermalization by iron whiskers of diameter $\sim 0.02 \mu\text{m}$ and length $\sim 1 \text{mm}$, which possess a flat absorption curve over the wavelength range $\sim 0.2 \text{mm} \approx 2 \text{cm}$, with an average mass absorption coefficient of $1.7 \times 10^7 \text{cm}^2 \text{g}^{-2}$ and a smoothed-out cosmological mass density of $10^{-34} \text{g cm}^{-3}$, the effective 'last emission' photosphere is an Earth-centred spherical surface of radius $R \approx 200 \text{Mpc}$ (refs 2–4). Fluctuations in the density distribution of iron needles in the foreground at distances nearer than 200 mpc, which are inevitable, would naturally lead to a microwave extinction which varies in intensity across the sky, but which does not alter the spectrum, leading in turn to fluctuations in the measured microwave background temperature. Some of these fluctuations could arise from clouds confined to the Galaxy, others could be of extragalactic origin.

With a reasonable assumption of density fluctuations of the order of 1 part in 10,000 occurring over inter-cluster distance scales of $L \approx 20 \text{Mpc}$, the resulting temperature fluctuations can be shown to be $\Delta T/T \approx 5 \times 10^{-6}$, as observed.

A typical angular scale of fluctuations would be $\Delta\theta \approx L/R \approx 6^\circ$, but larger fluctuations up to 90° could, of course, arise less frequently from a few local clouds. The Big Bang explanation of the latest COBE result is thus not unique, and alternative thermalization models cannot be excluded.

N. C. Wickramasinghe

School of Mathematics,
University of Wales College Cardiff,
Senghenydd Road,
Cardiff CF2 4AG, UK

1. Silk, J. *Nature* **356**, 741–742 (1992).
2. Hoyle, F. & Wickramasinghe, N. C. *Astrophys. Space Sci.* **147**, 245–256 (1988).
3. Wickramasinghe, N. C., Wickramasinghe, A.N. & Hoyle, F. *Astrophys. Space Sci.* (in the press).
4. Wickramasinghe, N. C. *Astrophys. Space Sci.* (in the press).