

12. Knight, M. E. & Turner, G. F. *Proc. R. Soc. Lond. B* **271**, 675–680 (2004).
 13. Schluter, D. *Trends Ecol. Evol.* **16**, 372–380 (2001).
 Supplementary information accompanies this communication on Nature's website.
 Competing financial interests: declared none.

Oscillatory motion

Quantum whistling in superfluid helium-4

Fundamental considerations predict that macroscopic quantum systems such as superfluids and the electrons in superconductors will undergo oscillatory motion when forced through a small constriction. Here we induce these oscillations in superfluid helium-4 (⁴He) by pushing it through an array of nanometre-sized apertures. The oscillations, which are detected as an audible whistling sound, obey the so-called Josephson frequency relation and occur coherently among all the apertures. The discovery of this property in ⁴He at the relatively high temperature of 2 K (2,000 times higher than the temperature at which a related but different phenomenon occurs in ³He) may pave the way for a new class of practical rotation sensors of unprecedented precision.

The Josephson effects in superconductors have received attention both as an aid to scientific understanding and for their technological importance¹. Analogous effects, including Josephson oscillations, have been observed^{2,3} in superfluid ³He below 1 mK. However, detection of oscillations at the Josephson frequency in superfluid ⁴He has remained elusive until now, despite almost four decades of attempts⁴.

Superconductors and superfluids are both described by a macroscopic wave function that includes amplitude and phase, ϕ . A chemical-potential difference, $\Delta\mu = \mu_2 - \mu_1$, between two baths of superfluid separated by an aperture causes the phase difference, $\Delta\phi = \phi_2 - \phi_1$, to change in accordance with the Josephson–Anderson phase-evolution equation

$$\frac{d\Delta\phi}{dt} = \frac{-\Delta\mu}{\hbar}$$

where \hbar is Planck's constant (h) divided by 2π and where $\Delta\mu/m_4 = \Delta P/\rho - S\Delta T$ (and m_4 is the mass of the ⁴He atom, ΔP is the pressure difference, ρ is the mass density, S is the entropy per unit mass, and ΔT is the temperature difference). A non-zero $\Delta\phi$ results in a superfluid current, $I(\Delta\phi)$, through the aperture. If $I(\Delta\phi)$ is periodic for 2π , a constant $\Delta\mu$ causes current to oscillate through the aperture at the Josephson frequency $f_J = \Delta\mu/h$. The periodicity in $I(\Delta\phi)$ can occur if the aperture acts like an ideal weak link^{3,5}, in which case $I(\Delta\phi) \propto \sin(\Delta\phi)$, or by the generation of 2π phase slips⁶, in which case $I(\Delta\phi)$

is expected to follow a sawtooth waveform.

The experimental set-up is shown in Fig. 1a (for methods, see supplementary information). We used an electrostatically driven diaphragm² to apply an initial pressure step between two baths of superfluid separated by an aperture array. The array consisted of 65×65 nominally 70-nm apertures spaced on a 3- μm square lattice in a 50-nm-thick silicon nitride membrane. After the pressure step, fluid flowed through the array and the chemical-potential difference relaxed to zero. When the output of a diaphragm position sensor, which monitored fluid flow, was connected to a set of headphones, we heard a clear whistling sound that passed from high to low frequency (audio recording in supplementary information).

By using Fourier transform methods, we extracted the frequency and amplitude of this whistle as a function of time throughout the transient. Immediately after the pressure step is applied, the temperatures on either side of the aperture array are equal and the entire $\Delta\mu$ is determined by the initial pressure head, ΔP_0 . Figure 1b shows that the initial frequency is proportional to the initial chemical-potential difference. The slope of the line agrees, within the systematic error of our pressure calibration, with the Josephson frequency formula ($f_J = m_4\Delta P_0/\rho h$).

Oscillations resulting from 2π phase slips are expected to have a velocity amplitude $\kappa/2l$, where $\kappa = h/m_4$ is the circulation quantum and l is an effective length for one aperture⁷. If, in addition, the oscillation in each of the N apertures occurs coherently, the amplitude of the diaphragm-displacement Fourier component at f_J is

$$X_0 = \alpha \frac{\rho_s N \kappa a}{4\pi f_J \rho A l}$$

where A is the area of the diaphragm, a is the area of a single aperture, and ρ_s is the superfluid density. The factor α would be $2/\pi$ for a sawtooth waveform, or unity for a sinusoid of the same peak amplitude. We find $\alpha \approx 0.6$, independent of temperature in the range where, if T_λ is the superfluid transition temperature, $T_\lambda - T$ is between 1.7 and 2.9 mK.

We conclude that the oscillation is a coherent phenomenon involving all the apertures in the array, and is possibly sawtooth in waveform. This coherence is remarkable, because earlier work using a single aperture showed that thermal fluctuations in the phase-slip nucleation process destroy time coherence in the rate of phase slippage, so that no Josephson oscillation exists⁸. However, it seems that thermal fluctuations are suppressed for an array — an observation that calls for further investigation⁹.

We have found that superfluid ⁴He in an array of small apertures behaves quantum coherently, oscillating at the Josephson frequency. Because these oscillations appear in ⁴He at a temperature 2,000 times higher than




Figure 1 Quantum oscillations in ⁴He. **a**, Experimental cell (see supplementary information for details). **b**, Whistle frequency plotted against the initial pressure, $\Delta P_0 = \rho\Delta\mu_0/m_4$. Temperature is in the range where, if T_λ is the superfluid transition temperature, $T_\lambda - T$ is 1.7–2.9 mK. A fit (solid line) to the data gives a slope of 78 Hz mPa⁻¹, with a systematic uncertainty of 20% arising from our pressure calibration. This agrees with the Josephson frequency relation $f_J = \Delta\mu/h$ value of 68.7 Hz mPa⁻¹. The oscillation is still present down to at least 150 mK below T_λ , where the healing length is much smaller than the aperture diameter and $I(\Delta\phi)$ is linear. The oscillation is presumably due to periodic 2π phase slips.

in superfluid ³He, it may be possible to build sensitive rotation sensors using much simpler technology than previously believed^{10–13}. This could find application in rotational seismology, geodesy and tests of general relativity.

E. Hoskinson, R. E. Packard, Thomas M. Haard

Physics Department, University of California, Berkeley, California 94720, USA
 e-mail: packard@socrates.berkeley.edu

1. Van Duzer, T. & Turner, C. W. *Superconductive Devices and Circuits* 2nd edn (Prentice-Hall, Upper Saddle River, 1999).
2. Pereversev, S. V., Loshak, A., Backhaus, S., Davis, J. C. & Packard, R. E. *Nature* **388**, 449–451 (1997).
3. Davis, J. C. & Packard, R. E. *Rev. Mod. Phys.* **74**, 741–773 (2002).
4. Tilley, D. R. & Tilley, J. in *Superfluidity and Superconductivity* 3rd edn, Ch. 7 (Institute of Physics, Bristol and Philadelphia, 1990).
5. Sukhatme, K., Mukharsky, Y., Chui, T. & Pearson, D. *Nature* **411**, 280–285 (2001).
6. Avenel, O. & Varoquaux, E. *Phys. Rev. Lett.* **55**, 2704–2707 (1985).
7. Packard, R. E. & Vitale, S. *Phys. Rev. B* **45**, 2512–2515 (1992).
8. Backhaus, S. & Packard, R. E. *Phys. Rev. Lett.* **81**, 1893–1896 (1998).
9. Chui, T., Holmes, W. & Penanen, K. *Phys. Rev. Lett.* **90**, 085301 (2003).
10. Packard, R. E. & Vitale, S. *Phys. Rev. B* **46**, 3540–3549 (1992).
11. Avenel, O., Hakonen, P. & Varoquaux, E. *Phys. Rev. Lett.* **78**, 3602–3605 (1997).
12. Schwab, K., Bruckner, N. & Packard, R. E. *Nature* **386**, 585–587 (1997).
13. Simmonds, R. W., Marchenkov, A., Hoskinson, E., Davis, J. C. & Packard, R. E. *Nature* **412**, 55–58 (2001).

Supplementary information accompanies this communication on Nature's website.

Competing financial interests: declared none.

brief communications arising online

♦ www.nature.com/bca

Marine ecology: Different measures of biodiversity
 J. R. Dolan (doi:10.1038/nature03320)

Reply: X. Irigoien, J. Huisman, R. P. Harris
 (doi:10.1038/nature03321)