

SHORT COMMUNICATION

Photosynthetic temperature adaptation during niche diversification of the thermophilic cyanobacterium *Synechococcus* A/B clade

Deana Pedersen and Scott R Miller

Division of Biological Sciences, The University of Montana, Missoula, MT, USA

We take an *in vivo* fluorescence approach to investigate photosynthetic adaptation by ecologically divergent members of the A/B clade of the hot spring cyanobacterium *Synechococcus*, the most thermotolerant of which defines the upper thermal limit for photosynthesis. During *Synechococcus* diversification, both photosystem II and the light-harvesting phycobilisome have evolved greater thermostability as the group has invaded higher temperature habitats, particularly for the most thermotolerant lineage. This enhanced function at higher temperatures has come at the cost of reduced performance at lower temperatures, and these trade-offs contribute to niche specialization in the clade. Molecular evolutionary analyses revealed specific adaptive protein changes in the most thermotolerant lineage. Our study advances our understanding of the origins of *Synechococcus* diversity.

The ISME Journal (2017) 11, 1053–1057; doi:10.1038/ismej.2016.173; published online 16 December 2016

Introduction

Cyanobacteria are the dominant primary producers in alkaline hot springs at temperatures below ca. 73 °C, the upper limit for photosynthetic life (Ward *et al.*, 2012). In these environments, members of the *Synechococcus* A/B clade exhibit markedly divergent thermal ecologies (Miller and Castenholz, 2000; Allewalt *et al.*, 2006), with more thermotolerant lineages in the group descended from a moderately thermophilic ancestor (Miller and Castenholz, 2000; Miller *et al.*, 2013). The *Synechococcus* radiation therefore makes an excellent system for investigating how microorganisms have diverged during adaptation to novel environments. For example, although thermostability of the carbon-fixing enzyme RuBisCO was not under selection during much of the history of *Synechococcus* diversification, an evolved increase in stability was essential for the invasion of the highest temperature habitats that support photoautotrophy (Miller *et al.*, 2013).

Temperature has a major effect on photosynthesis, and photosystem II (PSII), the multisubunit complex that extracts electrons from water at its oxygen-evolving complex (OEC) by light-driven redox chemistry, is the most temperature sensitive component of the photosynthetic apparatus (Berry and Björkman, 1980). Temperature also impacts the ability to efficiently harvest

solar energy. In most cyanobacteria, light is primarily harvested for photosynthesis by phycobilisomes (PBS), supramolecular, phycobiliprotein-containing antennae that funnel excitation energy to the photosystem reaction centers (Glazer, 1985). High temperature stress can result in the reversible dissociation and energetic uncoupling of PBS from a photosystem or, under severe stress, irreversible PBS denaturation (Inoue *et al.*, 2000). To better understand how *Synechococcus* has diverged during adaptation to different thermal environments, we have investigated the temperature dependence of PSII and PBS stability and function with an *in vivo* fluorescence approach.

Materials and methods

Temperature dependence of fluorescence

Strains were grown under a 12/12 h light–dark cycle at 55 °C as in Miller and Castenholz (2000). We monitored fluorescence emission at 680 nm of cell suspensions containing 1 µg ± 0.04 (s.e.) Chl *a* per ml with a Photon Technology International model QM-7/2005 (Lawrenceville, NJ, USA) rapid temperature change spectrofluorometer equipped with a Quantum Northwest TLC 50 thermoelectric temperature-controlled cuvette holder. For each sample, cells were excited with either 437 or 605 nm light to excite either PSII or PBS, respectively. Fluorescence was measured beginning at 40 °C and then over 2.5 °C increments as the sample was heated until the temperature at which either complete inhibition of PSII (437 nm) or PBS denaturing (605 nm), respectively, was reached. Cells were stirred with a magnetic stir bar, and the

Correspondence: SR Miller, Division of Biological Sciences, The University of Montana, 32 Campus Drive No. 4824, Missoula, MT 59812, USA.

E-mail: scott.miller@umontana.edu

Received 11 April 2016; revised 31 August 2016; accepted 31 October 2016; published online 16 December 2016




Figure 1 (a) Bayesian phylogeny of the 16S rRNA gene for strains in the *Synechococcus* A/B clade and outgroup taxon *Synechococcus* strain C9. Topology and branch lengths are from Ward *et al.* (2012). The scale bar is 0.05 nucleotide substitutions per site. All nodes in the tree have posterior probability support of 100%. (b) Temperature dependence of relative *in vivo* fluorescence monitored at 680 nm for *Synechococcus* strains excited at 437 nm. Strain colors are as in (a). Error bars are s.e. A relative fluorescence value of 1.0 corresponds to the following mean absolute fluorescence values (photons per s) \pm s.e.: 96 871 \pm 14 070.0 (C9); 23 575 \pm 1892.1 (OSB'); 42 808 \pm 3975.8 (OH20); 43 869 \pm 3975.8 (OH2); 21 692 \pm 690.3 (OH28). (c) Temperature dependence of relative *in vivo* fluorescence monitored at 680 nm for *Synechococcus* strains excited at 605 nm. Strain colors are as in (a). Error bars are s.e. A relative fluorescence value of 1.0 corresponds to the following mean absolute fluorescence values (photons per s) \pm s.e.: 315 815 \pm 19 215.7 (C9); 108 273 \pm 6220.4 (OSB'); 226 643 \pm 16 087.7 (OH20); 338 726 \pm 16 340.4 (OH2); 28 335 \pm 432.1 (OH28).

temperature in the cuvette was independently monitored with an Omega temperature probe for all measurements. Experiments were conducted in triplicate for independent cultures of each strain.

Correlation analyses of thermotolerance traits

To estimate Pearson's correlation coefficients between thermotolerance measures, we used phylogenetic generalized least-squares models (Martins and Hansen, 1997) implemented in Compare version 4.6 (<http://compare.bio.indiana.edu/>), which account for the statistical dependence of phenotypic data collected from related organisms by converting phylogeny branch lengths into units of expected phenotypic divergence. Branch length data were from the phylogeny in Ward *et al.* (2012; Figure 1a).

Comparative sequence analysis

Genomic DNA was isolated as in Miller and Castenholz (2000), and fragments of genes encoding components of PSII and the PBS were amplified using primers (Supplementary Table S1) designed from the *Synechococcus* strains A and OSB' genomes (GenBank accession nos CP000239 and CP000240; Bhaya *et al.*, 2007). These were directly sequenced on an ABI 3130 at the University of Montana DNA Sequencing Facility (Missoula, MT, USA). Sequence data were deposited in GenBank (accession numbers KX758263–KX758291). For each gene, we implemented two maximum-likelihood models of codon evolution with the program codeml in PAML version 3.14: a branch-site model (Yang and Nielsen, 2002) that tests for adaptive evolution at the subset of codon sites along a specified lineage by allowing ω (that is, d_N/d_S) to vary both among codon sites and among branches; and a null model that constrains ω between 0 and 1 (that is, neutral evolution). Models were compared with likelihood ratio tests. For analyses for which the null model could be rejected, positively selected codons were identified by Empirical Bayes analysis (probability cutoff of 90%).

Results and discussion

Strains from the *Synechococcus* A/B clade (Figure 1a) and the thermophilic outgroup strain *Synechococcus* C9, which had been isolated from different thermal niches along the environmental gradients of North American alkaline hot springs, have diverged in their temperature dependence of growth (Miller and Castenholz, 2000; Allewalt *et al.*, 2006; Table 1). These differences have persisted during laboratory maintenance: less thermotolerant strains C9 and OSB' did not grow at or above 61 °C, strain OH20 grew at 61 °C but not at 64 °C, strain OH2 grew at 64 °C and only OH28 was capable of growth at or above 65 °C. We used fluorescence signals of divergent *Synechococcus* strains grown at a common garden temperature of 55 °C (Table 1) to test whether there have been evolutionary changes in the temperature dependence of PSII and PBS function during the diversification of this group.

Table 1 *Synechococcus* strain growth at 55 °C and temperature dependence

Strain	55 °C growth rate (doublings day) \pm s.e.	T_{opt} (°C)	T_{max} (°C)
C9	0.79 \pm 0.008	45	55
OSB'	0.56 \pm 0.059	50	60
OH20	0.76 \pm 0.039	57	62
OH2	0.57 \pm 0.017	57	63
OH28	0.37 \pm 0.020	65	70

Abbreviations: T_{opt} , observed temperature of maximum growth rate values from Miller and Castenholz (2000) and Allewalt *et al.* (2006); T_{max} , observed maximum temperature supporting growth from Miller and Castenholz (2000) and Allewalt *et al.* (2006).

Because fluorescence and photochemical energy capture are competing de-excitation processes following light absorption, heat-induced impairment of PSII function results in an increased fluorescence yield (Berry and Björkman, 1980) as electron transport is inhibited between primary and secondary electron acceptors Q_A and Q_B (Cao and Govindjee, 1990). The temperature dependence of fluorescence emission during the excitation of PSII with blue light varied among divergent *Synechococcus* A/B strains and *Synechococcus* strain C9 (Figure 1b). Differences among strains explain 98% of the variation in the temperature at which the fluorescence maximum is reached (that is, the temperature of PSII inactivation; $P < 0.0001$), which indicates that the onset of heat-induced damage has evolved during *Synechococcus* diversification. Based on *t*-tests with a Bonferroni correction to account for multiple pairwise comparisons, the temperature of PSII inactivation increased in the order: C9 < OSB' < OH20 = OH2 < OH28 ($P \leq 0.005$ for all significant comparisons). This is most striking for *Synechococcus* strain OH28, for which PSII function was not compromised at temperatures sufficient to inactivate all PSII centers of less thermotolerant strains. This strain was also characterized by lower baseline levels of relative chlorophyll fluorescence compared with the other strains (Figure 1b). In addition to photochemical quenching, other relaxation processes that compete with fluorescence following chlorophyll excitation that may potentially differ between OH28 and other strains include energy transfer to other pigments (including PSI), heat dissipation and the distribution of chlorophyll between PSII and PSI. Whether any of these fluorescence-quenching processes contribute to these differences, as well as whether there is a mechanistic connection to the extreme temperature tolerance of PSII in strain OH28, is a subject for future investigations. After accounting for the phylogenetic dependence of the data (see Materials and methods), the temperature of PSII inactivation was strongly positively correlated with both strain optimal growth temperature ($R = 0.98$) and maximum growth temperature ($R = 0.98$) measured in previous studies (Table 1; Miller and Castenholz, 2000; Allewalt *et al.*, 2006). The effect of short-term heat

stress on PSII function is therefore associated with the long-term temperature dependence of strain fitness.

We next assayed the thermostability of *Synechococcus* PBS. When PBS are excited at 605 nm, fluorescence emission monitored at 680 nm has two major sources: PSII and the PBS itself (Campbell *et al.*, 1998). Fluorescence is at a minimum when both photochemical efficiency and energetic coupling between PBS and PSII are high (Pospíšil *et al.*, 1998; Tamary *et al.*, 2012). With further heating, fluorescence initially increases to a local maximum due to both decreased photochemical efficiency and heat-induced energetic uncoupling of membrane-diffusible PBS from the photosystems, but then sharply declines as PBS irreversibly denature (Inoue *et al.*, 2000). As we observed above for PSII, there has also been substantial evolution of PBS thermostability during *Synechococcus* diversification, most notably for strain OH28 (Figure 1c). Differences among strains in the measured temperature of the fluorescence maximum under heating were very highly significant overall ($P < 0.0001$), ranged from 62.5 °C (strain C9) to 80 °C (strain OH28), and increased in the order: C9 < OSB' = OH20 = OH2 < OH28 ($P \leq 0.005$ for all significant *t*-test comparisons with a Bonferroni correction). The temperature of this fluorescence maximum is strongly positively correlated with both optimal ($R = 0.96$) and maximum ($R = 0.99$) growth temperatures of *Synechococcus* (Table 1). Although the fluorescence maximum was not significantly different for strains OSB', OH20 and OH2, these strains were distinguished by subtle but statistically significant differences in PBS stability based on the dynamics of the steep fluorescence decrease that is indicative of phycobiliprotein denaturation (Figure 1c; $P < 0.02$ for the comparison between OH20 and OSB' at 72.5 °C and $P < 0.001$ for comparisons between OH2 and both OH20 and OSB' at 75 °C). Strains with higher temperature maxima for growth (Table 1), therefore, have more stable PBS. In addition, the position of the fluorescence minimum, where photochemical quenching is greatest, is observed at higher temperatures in strain OH28 compared with other strains (Figure 1c), indicating that enhanced function at higher temperatures comes at the cost of reduced photochemical efficiency at lower temperatures. The inability of the photosynthetic apparatus to function well at both high and low temperatures thus contributes to the niche specialization of *Synechococcus*.

Taken together, the above results emphasize that enhanced thermostability of both PSII and PBS has contributed to the adaptation to higher temperatures during diversification of the *Synechococcus* A/B clade, particularly in the most thermotolerant lineage represented by strain OH28. To gain insights on the molecular basis of these changes, we implemented maximum-likelihood models of codon evolution (Yang and Bielawski, 2000) for several PSII and PBS

genes from these strains. These included genes from the PSII core (*psbA*, *psbB*, *psbD*) and *psbO*, which encodes an extrinsic protein that forms a cap over the oxygen-evolving complex, as well as genes contributing to the membrane-associated core (*apcA*, *apcB*, *apcD*, *apcE*) and distal rods (*cpcA*, *cpcB*) of the PBS (Supplementary Table S1). Owing to the degeneracy of the genetic code, a nucleotide substitution in a protein-coding gene may or may not alter the amino-acid sequence. These models estimate the ratio of nonsynonymous (that is, amino-acid changing; d_N) to synonymous (d_S) substitution rates ($\omega = d_N/d_S$). Nonsynonymous mutations are typically expected to be deleterious and therefore selected against ($\omega < 1$), whereas an excess of nonsynonymous substitutions ($\omega > 1$) is indicative of adaptive evolution (Yang and Bielawski, 2000). For each locus, we implemented a branch-site model (Yang and Nielsen, 2002) that tested for adaptive evolution at a few key sites along each branch of the *Synechococcus* phylogeny by allowing ω to vary both among codon sites and among branches (see Materials and methods). These models can be statistically compared by a likelihood ratio test with a null model for which ω is constrained between 0 and 1.

Amino-acid changes were very rare overall for all genes ($\omega < 0.1$). Despite these functional constraints on protein evolution, two genes exhibited evidence for a class of codon sites that was under positive selection along the strain OH28 terminal branch of the *Synechococcus* phylogeny presented in Figure 1a. The first was *psbA*, which encodes the D1 protein. D1 contributes most of the residues that bind and stabilize the oxygen-evolving complex, the reaction center and other electron transport components on both the donor and acceptor sides of PSII (Ferreira *et al.*, 2004). One site (codon 222) was assigned to the positively selected class ($\omega = 4.52$; $2\Delta L = 16.76$, $P = 0.0002$) with high probability (97%). This site, which is threonine in strain OH28 but a highly conserved serine in other publicly available *psbA* sequences, is near the binding pocket of the secondary electron acceptor Q_B . This is of potential interest, given that inhibition of electron transport between Q_A and Q_B is likely responsible for heat-induced PSII inactivation (Cao and Govindjee, 1990). The other positively selected gene in our analyses was *cpcA* ($\omega = 3.17$; $2\Delta L = 7.56$, $P = 0.025$), which encodes the α -subunit of the PBS rod phycobiliprotein phycocyanin. One site (codon 111), at which glutamine has been replaced by methionine in strain OH28, was inferred to belong to the positively selected site class (90%). This residue makes contacts with the $\alpha 84$ phycocyanobilin chromophore and may contribute to changes in the protein environment surrounding the chromophores that are responsible for the observed blue shift in the absorption maximum of phycocyanin from ~ 620 to 608 nm in strain OH28 (unpublished data) and a related strain (Edwards *et al.*, 1996).

These results also inform our understanding of the impacts of acute thermal stress at the physiological limits of photosynthetic metabolism. Whereas, in most strains, PSII is inactivated at approximately the same temperature as that of PBS dissociation from the photosystems (Figure 1), PSII of strain OH28 is more thermostable than the PBS. That is, the phenotypic effects of adaptive amino-acid substitutions that have occurred along the OH28 lineage (either positively selected substitutions detected by our models of codon evolution and/or unobserved changes at other genetic loci) have had a larger impact on the temperature dependence of PSII function, resulting in an asymmetric response exclusive to this lineage during *Synechococcus* diversification. Although the temperatures required to inactivate PSII and PBS exceed the thermal limit for sustained growth of each strain (Table 1; Miller and Castenholz, 2000; Allewalt *et al.*, 2006), this shift is nonetheless significant, because it changes the nature of the physiological response of the photosynthetic apparatus during short-term exposure to extreme but non-lethal thermal stress that may occur in geothermal environments.

Conflict of Interest

The authors declare no conflict of interest.

Acknowledgements

We thank two anonymous reviewers for their comments and suggestions on an earlier version of the paper. We also thank Reid Longley for his technical assistance with the fluorescence experiments and Big Sky High School (Missoula, MT, USA) interns Amanda Tripp and Amy Smith, who learned the polymerase chain reaction as a part of this project. This work was supported by US National Science Foundation award MCB-0347627 and NASA Astrobiology Institute award NNA15BB04A.

References

- Allewalt JP, Bateson MM, Revsbech NP, Stack K, Ward DM. (2006). Effect of temperature and light on growth of and photosynthesis by *Synechococcus* isolates typical of those predominating in the Octopus Spring microbial mat community of Yellowstone National Park. *Appl Environ Microbiol* **72**: 544–550.
- Berry J, Björkman O. (1980). Photosynthetic response and adaptation to temperature in higher plants. *Ann Rev Plant Physiol* **31**: 491–543.
- Bhaya D, Grossman AR, Steunou AS, Khuri N, Cohan FM, Hamamura N *et al.* (2007). Population level functional diversity in a microbial community revealed by comparative genomic and metagenomic analyses. *ISME J* **1**: 703–713.
- Campbell D, Hurry V, Clarke AK, Gustafsson P, Oquist G. (1998). Chlorophyll fluorescence analysis of cyanobacterial photosynthesis and acclimation. *Microbiol Mol Biol Rev* **62**: 667–683.

- Cao J, Govindjee. (1990). Chlorophyll *a* fluorescence transient as an indicator of active and inactive photosystem II in thylakoid membranes. *Biochim Biophys Acta* **105**: 180–188.
- Edwards MR, MacColl R, Eisele LE. (1996). Some physical properties of an unusual C-phycoyanin isolated from a photosynthetic thermophile. *Biochim Biophys Acta* **1276**: 64–70.
- Ferreira KN, Iverson TM, Maghlaoui K, Barber J, Iwata S. (2004). Architecture of the photosynthetic oxygen-evolving center. *Science* **303**: 1831–1838.
- Glazer AN. (1985). Light-harvesting by phycobilisomes. *Annu Rev Biophys Chem* **14**: 47–77.
- Inoue N, Emi T, Yamane Y, Kashino Y, Koike H, Satoh K. (2000). Effects of high-temperature treatments on a thermophilic cyanobacterium *Synechococcus vulcanus*. *Plant Cell Physiol* **41**: 515–522.
- Martins EP, Hansen TF. (1997). Phylogenies and the comparative method: a general approach to incorporating phylogenetic information into the analysis of interspecific data. *Am Nat* **149**: 646–667.
- Miller SR, Castenholz RW. (2000). Evolution of thermotolerance in hot spring cyanobacteria of the genus *Synechococcus*. *Appl Environ Microbiol* **66**: 4222–4229.
- Miller SR, McGuirl MA, Carvey D. (2013). The evolution of RuBisCO stability at the thermal limit of photoautotrophy. *Mol Biol Evol* **30**: 752–760.
- Pospišil P, Skotnica J, Nauš J. (1998). Low and high temperature dependence of minimum F_0 and maximum F_M chlorophyll fluorescence *in vivo*. *Biochim Biophys Acta* **163**: 95–99.
- Tamary E, Kiss V, Nevo R, Adam Z, Bernát G, Rexroth S *et al.* (2012). Structural and functional alterations of cyanobacterial phycobilisomes induced by high-light stress. *Biochim Biophys Acta* **1817**: 319–327.
- Ward DM, Castenholz RW, Miller SR. (2012). Cyanobacteria in geothermal habitats. In: Whitton BE (ed). *Ecology of Cyanobacteria II*. Springer: Dordrecht, The Netherlands.
- Yang Z, Bielawski J. (2000). Statistical methods for detecting molecular adaptation. *Trends Ecol Evol* **15**: 496–503.
- Yang Z, Nielsen R. (2002). Codon-substitution models for detecting molecular adaptation at individual sites along specific lineages. *Mol Biol Evol* **19**: 908–917.

Supplementary Information accompanies this paper on The ISME Journal website (<http://www.nature.com/ismej>)