

OPEN
ARTICLE

A curated collection of human vaccination response signatures

Kenneth C. Smith^{1,9}, Daniel G. Chawla^{2,9}, Bhavjinder K. Dhillon^{3,9}, Zhou Ji¹, Randi Vita⁴, Eva C. van der Leest³, Jing Yi Jessica Weng³, Ernest Tang³, Amani Abid³, The Human Immunology Project Consortium (HIPC)*, Bjoern Peters^{4,5}, Robert E. W. Hancock^{3,10}, Aris Floratos^{1,6,10} & Steven H. Kleinstein^{2,7,8,10}

Recent advances in high-throughput experiments and systems biology approaches have resulted in hundreds of publications identifying “immune signatures”. Unfortunately, these are often described within text, figures, or tables in a format not amenable to computational processing, thus severely hampering our ability to fully exploit this information. Here we present a data model to represent immune signatures, along with the Human Immunology Project Consortium (HIPC) Dashboard (www.hipc-dashboard.org), a web-enabled application to facilitate signature access and querying. The data model captures the biological response components (e.g., genes, proteins, cell types or metabolites) and metadata describing the context under which the signature was identified using standardized terms from established resources (e.g., HGNC, Protein Ontology, Cell Ontology). We have manually curated a collection of >600 immune signatures from >60 published studies profiling human vaccination responses for the current release. The system will aid in building a broader understanding of the human immune response to stimuli by enabling researchers to easily access and interrogate published immune signatures.

Introduction

Systems-level profiling of the human immune system has generated important insights into the mechanisms by which humans respond to exposures such as vaccination. These studies, including many conducted through the Human Immune Project Consortium (HIPC), have generated hundreds of publications. While repositories exist to promote re-use of primary experimental immunology data generated from these efforts, such as the Gene Expression Omnibus¹ (GEO) and the NIAID Division of Allergy, Immunology, and Transplantation (DAIT)-sponsored Immunology Database and Analysis Portal² (ImmPort), there is no centralized framework to aggregate and organize the published findings resulting from the analysis of this data, and particularly the coherent sets of biomarkers, termed here “signatures”. Additionally, such signatures are not published in a consistent format between publications and may be presented as text, tables, or images. This heterogeneity presents a barrier to comparative analyses since identifying published signatures, for example of a vaccine response, requires extensive manual curation of the literature that must be repeated by investigators each time they wish to interpret a set of results. Here, we propose a model to standardize the representation of these published findings and present the Human Immunology Project Consortium (HIPC) Dashboard—a searchable interface to query curated signatures from the corpus of human immunology literature.

We define a ‘signature’ as the information required to specify a published result. This includes alterations in the levels of a set of one or more response component(s), i.e., biological entities such as genes or cell types, that are

¹Department of Systems Biology, Columbia University, New York, New York, USA. ²Program in Computational Biology and Bioinformatics, Yale University, New Haven, CT, USA. ³Department of Microbiology & Immunology, University of British Columbia, Vancouver, Canada. ⁴Center for Infectious Disease and Vaccine Research, La Jolla Institute for Immunology, La Jolla, CA, USA. ⁵Department of Medicine, Division of Infectious Diseases and Global Public Health, University of California, San Diego, La Jolla, CA, USA. ⁶Department of Biomedical Informatics, Columbia University, New York, New York, USA. ⁷Department of Pathology, Yale School of Medicine, New Haven, CT, USA. ⁸Department of Immunobiology, Yale School of Medicine, New Haven, CT, USA. ⁹These authors contributed equally: Kenneth C. Smith, Daniel G. Chawla, Bhavjinder K. Dhillon. ¹⁰These authors jointly supervised this work: Robert E. W. Hancock, Aris Floratos, Steven H. Kleinstein. *A list of authors and their affiliations appears at the end of the paper. ✉e-mail: af2202@cumc.columbia.edu; steven.kleinstein@yale.edu

defined by a particular comparison in the context of an immune exposure. The signature also includes contextual information (termed metadata) such as the conditions and circumstances under which the signature was identified, the tissues or cells that were assayed, as well as clinical data such as demographic information about the groups that were included in the analysis. As a motivating example, a study by Bucasas *et al.*³ identified a set of genes that are up-regulated in individuals with higher antibody responses (comparison) after vaccination with the 2008–2009 trivalent influenza vaccine (exposure) in an adult cohort. The expression of genes STAT1, IRF9, SPI1, CD74, HLA-E, and TNFSF13B one day after influenza vaccination was predictive of greater antibody responses. In the paper, these results were represented in a table, though similar findings often appear as text or within figures. Without standardization, such findings are not easily accessible to the wider scientific community for further analysis.

Several existing resources define pathway and gene module signatures through re-analysis of raw data, but few capture the original findings published with these data or are specifically geared towards human immunology research. Among these resources are the OMICS Compendia Commons (OMiCC)⁴, EnrichR^{5,6}, the integrative Library of Integrated Network-based Cellular Signatures (iLINCS)⁷, the Molecular Signatures Database (MSigDB)^{8,9}, and VaximmutorDB¹⁰. OMiCC crowdsources annotations for gene expression data to be used in re-analysis and novel signature generation. EnrichR and iLINCS offer biological annotations built from data re-analyzed en masse, but similarly do not capture published findings. MSigDB does include manually curated gene signatures along with those derived from data re-analysis, albeit with fewer contextual details than captured for the HIPC Dashboard. VaximmutorDB captures published gene expression and proteomic signatures but not cell-type frequency signatures, and signatures from this database are not yet downloadable in a machine-readable format.

To improve access and to promote reuse of published signatures, we designed a data model that standardizes the content and context of published immune signatures. Our initial curation efforts have focused on gene expression and cell-type frequency/activation signatures of human vaccine responses, but this framework is extensible to other domains such as response to infection. We captured what is changing, (e.g. groups of genes), how that response component changed (e.g. up- or down-regulation), where this change was observed (e.g. in sorted CD8+ T cells from adults), and the comparison that was performed (e.g. individuals with high vs. low antibody titers post-vaccination). We then manually curated signatures from publications both within and outside of HIPC that described changes in gene expression, cell-type frequencies, or cell activation state in response to vaccination. To disseminate these immune signatures, we developed the HIPC Dashboard (www.hipc-dashboard.org), a web-accessible, user-friendly interface to enable signature searching and browsing, and to facilitate rapid comparative analyses. The design of the HIPC Dashboard is based on a similar infrastructure we developed previously for the Cancer Target Discovery and Development network, the CTD² Dashboard¹¹, and leverages the same underlying ontological framework for the standardized representation of research findings as well as the emphasis on the consistent, curation-mediated use of controlled vocabularies for linking findings reported in different publications.

Results

A data model for immune signatures of vaccination. We developed a data model that captures, in a detailed and consistent format, the essential information embedded in published immune signatures of vaccination for dissemination through the HIPC Dashboard (Table 1). Key elements of this data model (e.g., genes, vaccines, etc.) are specified using controlled vocabularies, thus making immune signatures of vaccination amenable to data mining and promoting compatibility with projects both within and outside of HIPC. A signature as defined in this model encapsulates both a change in the behavior or abundance of a biological response component as well as the metadata describing the context under which the signature is identified, including (1) the tissue in which the signature was observed, (2) the immune exposure and timing underlying the observed comparison, and (3) clinical details of the cohort from which tissue samples were taken, including age (Fig. 1). The model accommodates many types of biological response components (gene, protein, metabolite, pathway, and cell type (e.g. subsets of blood cells)). We focused on gene expression and cell type signatures of vaccination, but the data model and HIPC Dashboard infrastructure are flexible and can be easily expanded to accommodate arbitrary signature types.

To facilitate data mining and comparative analyses between different conditions including vaccine types, and to afford consistency between this database and other projects using the same controlled vocabulary terms, standardized terms and ontology links were used for as many biological response components, immune exposures, and demographic fields as possible. Gene and cell type response components were standardized to the HGNC¹² (as provided through the NCBI) and Cell Ontology^{13,14} (CL) respectively to enable comparisons across publications using different naming conventions. Cell types from the Cell Ontology were further differentiated using protein marker terms drawn from Protein Ontology¹⁵ (PRO), where possible (e.g. IFNG+ T cells). This same naming convention is used to describe the tissue in which the signature was observed¹⁶.

To annotate the immune challenges driving each signature, we utilized the Immune Exposure model¹⁷, which provides a standardized description of a broad range of potential and actual exposures to different immunological agents (e.g., vaccination, laboratory confirmed infection, living in an endemic area, etc.). Immune exposures are broken down into Exposure Process, Exposure Material, Disease Name, and Disease Stage. Each of these components is modeled using standardized ontology terminology. Within the data model for the HIPC Dashboard, Exposure Materials such as vaccines are captured using terms in the Vaccine Ontology¹⁸ (VO), which further link to target pathogens and strains using the NCBI Taxonomy^{19,20}. While these ontology choices reflect our initial focus on vaccination, the data model can accommodate other exposure processes beyond vaccination, with links to appropriate ontologies. Integration of the Immune Exposure model in the HIPC Dashboard data model promotes interoperability with other projects that have adopted its use, both within and outside of HIPC, including data repositories such as ImmPort and the Immune Epitope Database²¹.

Elements	Content Definition	Field value Example	Ontology	Data Type	Content Format
Response Component	The biological entity being observed: either a specific gene or cell subset	CKS1B	NCBI/HGNC for genes, Cell Ontology for cell types, PRO for proteins)	String	Controlled
Response Component Type	Type of response agent, e.g. gene, cell subset	gene		String	Fixed List
Tissue Type	The type of cells analyzed, e.g. whole blood, gated cells	blood (UBERON:0000178)	CL/PRO	String	Controlled
Exposure Process Type *	Category of immune exposure	Vaccination		String	Fixed List
Exposure Material *	Eg, the type of vaccine administered	VO:0001176	VO	String	Controlled
Disease Name *	The condition being observed	None	DO	String	Controlled
Disease Stage *	Stage of infection (e.g. acute, chronic, post, etc.)	None	OGMS	String	Controlled
Target Pathogen	The pathogenic organism (e.g. virus, bacterium, prion, fungus) being studied	Influenza A, Influenza B	NCBI Taxonomy ID	String	Controlled
Vaccine Year	Year for seasonal vaccines	2012		Numeric	Free Text
Adjuvant	A substance added to vaccines to increase the body's immune response to the vaccine		VO	String	Controlled
Route	Eg oral, nasal spray, intradermal injection, etc.	Intramuscular, intradermal, transcutaneous	VO	String	Controlled
Scheduling	The number of times a substance is administered within a specific time period.	e.g. prime / boost scheme		String	Free Text
Time Point	Time point relative to baseline event at which response was measured	1		Numeric	
Time Point Units	{hours, days, months, years}	day		String	Fixed List
Baseline Time Event	The starting point against which other events are compared	time of vaccination (first dose)		String	Free Text
Cohort	Features that describe study cohorts	e.g. antibody responders		String	Free Text
Age Min	Age of youngest subject	18		Numeric	
Age Max	Age of oldest subject	45		Numeric	
Age Units	{days, weeks, months, years}	years		String	Fixed List
Publication Reference	PubMed unique identifier of an article.	30843873		Numeric	Controlled
Publication Year	The year in which the study was published	2019		Numeric	
Publication Reference URL	Link to article in PubMed	https://www.ncbi.nlm.nih.gov/pubmed/30843873		String	URL
Comparison	The contrast used for deriving the signature	1d-0d		String	Free Text
Response Behavior	Observed change in the response agent under the comparison	Up		String	Free Text

Table 1. Data model for capturing immune signatures. Genes and cell types are captured as response components, with terms standardized against NCBI/HGNC or CL + PRO, respectively. Exposures are captured and standardized against VO terms and NCBI Taxonomy IDs. Metadata includes observed tissue, study timing, cohort descriptors, and age characteristics. *fields in the Immune Exposure model¹⁷.

Cohort information that is important for interpreting signatures is also captured. Cohort descriptors can vary widely between studies and can include, for example, sex, antibody response titers, geographic location, health status, vaccination or infection history, etc. This information is currently recorded as unstructured text to maintain flexibility. Cohort age range is standardized separately by storing minimum and maximum ages along with their units. Additional fields describe the particular perturbations that drive the changes to the biological response components. The “comparison” field describes the cohort groups whose differential response under the perturbation is measured. Examples of comparison groups include measurements taken at two different time points (e.g. day 1 vs. day 0), correlation with antibody response, differing antibody response outcomes (e.g. high vs. low responders), or comparisons across different demographic parameters such as age or sex (e.g. younger vs. older, female vs. male). The “response_behavior” field captures the directionality of the differential response (e.g. up or down, positively- or negatively-correlated) under the specified comparison. Fields for which a formal controlled vocabulary was not used, such as cohort descriptions and the comparison, are stored as free text.

Finally, each signature is tagged with a Pubmed ID (PMID) and publication year field, to connect observations to their literature sources.

Manual curation of published signatures. A defined Pubmed search strategy was used (see Methods) to assemble an initial list of publications comprising studies that involved a systems-level profiling of measurable changes before and after vaccination in human subjects. The publications were culled for signatures reporting statistically significant changes in gene expression, cell-type frequency or cell activation state induced by an immune exposure when comparing groups with different features (such as high vs low responders as defined by antibody titers). We focused on components that also included information about response behavior (e.g. up- or down-regulated). In total, 665 immune signatures were manually curated from 69 published studies. After standardization and quality control (see Methods), these curated gene and cell type signatures included 13,812 unique genes, 152 unique cell types (including protein markers and additional type-modifiers), and 44 pathogens across 56 vaccines (Table 2). Table 3 illustrates a typical gene-expression type signature after tissue, gene symbol and pathogen standardization.




Fig. 1 Overview of the manual curation process for extracting immune signatures from relevant publications into the HIPC signatures database and a web-accessible HIPC Dashboard. The middle panel highlights the various fields that are captured for a given immune signature, with examples provided in red font. Key fields are standardized using existing ontologies or pre-defined criteria in order to capture a wide array of signatures.

The HIPC Signatures Dashboard. The data model provides a means for representing immune signatures in a structured, standardized, and machine-readable manner while the curation process enables the cross-referencing of signatures from different publications based on their shared response components, by enforcing the consistent use of controlled vocabularies for codifying these components (e.g., genes, cell types, tissues, vaccines, and pathogen strains). These capabilities come together in the “HIPC Dashboard” (<http://hipc-dashboard.org>), a web application developed to enable dissemination of the curated set of immune response signatures. The HIPC Dashboard allows signature browsing, as well as searching for one or multiple response components (using the corresponding controlled vocabulary terms and their synonyms), to retrieve all immune signatures involving the query response component(s) across all curated publications.

The central viewable element of the HIPC Dashboard is the “Observation Summary”, a human-readable description of the information captured in an immune signature. Observation Summaries are constructed “on the fly”, using template text devised as part of the curation process. The template has placeholders for the various elements of an immune signature, including the response component (gene or cell type) and the response behavior type (up/down or correlation). When a specific signature is selected in the process of browsing or searching the Dashboard, the observation summary for that signature is instantiated by replacing the template placeholders with the relevant values from that signature. For example, a joint search on the terms “CD4” and “Zostavax” yields about 35 observation summaries. One of these is related to a change in cell-type frequency:

In peripheral blood mononuclear cell, CD4-positive, alpha-beta memory T cell & CD38+, HLA-DR+, VZV tetramer+ frequency was up at 14 days from time of vaccination for the comparison 14d vs 0d in cohort 50-75 yo after exposure to Zostavax targeting Human alphaherpesvirus 3 ([details »](#))

Here, the “&” sign separates the Cell Ontology cell type from Protein Ontology surface and other markers. In a second example, a joint search on the terms “CXCL10” and “BCG” yields 6 observation summaries, one of which reports a correlation of gene expression (at 1 day post-vaccination) to an ELISpot result at 28 days post-vaccination:

In blood, CXCL10 gene expression at 1 day from time of vaccination was positively correlated with IFN-gamma ELISpot spot forming cell 28d in cohort 4-6 mo subgroup BCG-primed after exposure to MVA85A targeting Mycobacterium tuberculosis variant bovis BCG ([details »](#))

Signature Type	Vaccines	Target Pathogens	Publications	Signatures	Total Response Components	Response Components per Signature
Gene	52	38	62	480	13,812 genes	1 to 2,036 genes
Cell type	28	26	31	185	47 cell types	1 to 9 cell types
Joint	56	44	69	665		

Table 2. Dashboard summary statistics for gene and cell-type signatures. “Joint” refers to the union of the two signature types, as they overlap in the various components. “Total Response Components” lists the number of genes or parent cell types from the Cell Ontology (CL) across all signatures. When additional cell-type markers are included, e.g. from the Protein Ontology (PRO), there are 152 unique cell types represented among the signatures. “Response Components per Signature” shows the range of the number of response components found in individual signatures.

Column name	Ontology	Values
response_component	HGNC	STAT1, IRF9, SPI1, CD74, HLA-E, TNFSF13B
tissue_type	Cell Ontology	CL:2000001 (peripheral blood mononuclear cell)
exposure_material	Vaccine Ontology	VO:0000045 (Fluarix); VO:0000046 (Fluvirin)
target_pathogen	NCBI Taxonomy	504904 (Influenza A virus (A/Brisbane/59/2007(H1N1))); 476294 (Influenza A virus (A/Brisbane/10/2007(H3N2))); 461739 (Influenza B virus (B/Florida/4/2006))
vaccine_year		2008
time_point		1
time_point_units		days
baseline_time_event		time of vaccination (first dose)
cohort		subgroup high responders
publication_reference (PMID)		21357945
comparison		correlated with titer response index (TRI)
response_behavior		positive

Table 3. Key fields in the immune signature data model for a gene expression signature. This signature reports positive correlation between gene expression and a computed titer response index (TRI).


In both cases, placeholders in the observation summary template have been replaced by controlled terms for the response components and ontology-linked metadata (blue, hyperlinked text) and by free-text metadata describing informative experiment details (black, bold text). Following the hyperlink for a controlled term leads to a dedicated page for the corresponding biological entity, providing additional details (including links to relevant external annotation sources, e.g., Entrez, GeneCards and UniProt for genes) as well as a listing of all the immune signatures stored in the Dashboard that involve that entity (Fig. 2a). Further, the “details” link at the end of each observation summary points to an “Observation” page (Fig. 2b) containing detailed information about the corresponding immune signature, including a full listing of all its available metadata. This includes, for example, structured text for values such as age group and days post-immunization, and links to download the full signature source data (including all metadata) in tab-delimited form. Additionally, each observation includes a link to a file containing the complete set of response components from which it was derived, e.g. the full list of genes or cell types.

Discussion

Users of the HIPC Dashboard can easily search and examine hundreds of immune signatures related to human vaccination responses. Consolidating these publications, standardizing their findings in a database, and disseminating them through the Dashboard interface allows for rapid comparative analyses and re-use of published findings. This is particularly important for identifying commonalities across studies that may reflect shared mechanisms. The HIPC Dashboard can offer broad insights into the mechanisms by which our immune systems respond to vaccination and will be of great value to the vaccine research community. Although the HIPC Dashboard is not designed as an analysis engine, all signatures are made available for download so that users may perform more sophisticated and targeted downstream analyses.

Among data resources dedicated to the collection of vaccination signatures, the HIPC Dashboard is nearly unique in its emphasis on manual curation of published literature. To the best of our knowledge, only MSigDB and VaximmutorDB maintain signatures curated from publications. MSigDB provides minimally redundant gene sets for enrichment analyses, but unlike the HIPC Dashboard does not attempt to capture the full biological context of published results. A reduced set of our curations has recently been made available for gene set enrichment analyses through MSigDB under the C7 VAX gene sets. VaximmutorDB provides access to a collection of immune factors (genes/proteins) that change in response to vaccination against 46 pathogens. Compared to VaximmutorDB, the HIPC Dashboard offers several advantages, including: (i) a wider breadth in the types of response components and immune changes that are captured, (ii) improved browsing functionality

a CD4-positive, alpha-beta T cell



Cell Ontology ID	CL_0000624
Definition	A mature alpha-beta T cell that expresses an alpha-beta T cell receptor and the CD4 coreceptor.
Comment	
Broad Synonyms	
Exact Synonyms	<ul style="list-style-type: none"> CD4-positive, alpha-beta T lymphocyte CD4-positive, alpha-beta T-cell CD4-positive, alpha-beta T4lymphocyte
Related Synonyms	
References	Cell Ontology: CL_0000624 EMBL-EBI OLS: CL_0000624

Related observations for the role of cell_biomarker

Show Connected Subjects Word Cloud

Showing 1 to 10 of 67 entries [Export as Spreadsheet](#) Filter this Table:

Publication Date	Observation Summary	PMID
Mar 16, 2019	In peripheral blood mononuclear cell, CD4-positive, alpha-beta T cell frequency was up at 0 days from time of vaccination for comparison females vs males in cohort 50-74 yo after exposure to Fluorix targeting Influenza A virus (A/California/7/2009(H1N1)), Influenza A virus (A/Perth/16/2009(H3N2)) and/or Influenza B virus (B/Brisbane/60/2008) (details »)	30873150
Sep 25, 2018	In blood, CD4-positive, alpha-beta T cell & IFNG+, VZV-specific frequency was up at 72 hours from time of vaccination for comparison 72hrs post-vaccination vs pre-vaccination in cohort 70-93 yo after exposure to Zostavax targeting Human alphaherpesvirus 3 (details »)	30247603

b Observation

In blood, CD4-positive, alpha-beta T cell & IFNG+, VZV-specific frequency was up at 72 hours from time of vaccination for comparison 72hrs post-vaccination vs pre-vaccination in cohort 70-93 yo after exposure to Zostavax targeting Human alphaherpesvirus 3

Name	Class	Role	Description
 blood	CellSubset	tissue	tissue (Cell Ontology)
 CD4-positive, alpha-beta T cell	CellSubset	cell_biomarker	response component
 Human alphaherpesvirus 3	Pathogen	pathogen	target pathogen (NCBI TaxID)
 Zostavax	Vaccine	vaccine	exposure material (vaccine ontology)

Study (show details)

Evidence

Showing 1 to 35 of 35 entries Filter this Table:

Type	Description	Details
------	-------------	---------

Fig. 2 HIPC Dashboard web interface. (a) Subject page for cell type “CD4-positive, alpha-beta T cell” showing a link-out to the Cell Ontology, the filtering box to further narrow the displayed observations, and the first two observation summaries (“Related observations”). (b) Partial view of a details page for a CD4-positive, alpha-beta T cell observation. For each controlled term, its name, plus its class, role, and description are shown. Linked pages list details from the relevant ontology and list all observations containing that term. The class equates to its controlled vocabulary type; values are cell subset, gene, pathogen, and vaccine. Roles are used to further differentiate how each term, whether controlled or standardized, is being used. Among the classes in the HIPC Dashboard, only the class “cell subset” has more than one role, these being “tissue” and “cell_biomarker”. Full metadata, not shown here, is contained in the table labeled “Evidence” at the bottom.

that facilitates comparisons of immune changes across studies and vaccines, and (iii) the ability to download signature data.

As of the date of this publication, 152 unique cell types and 13,812 distinct genes have been collected in the HIPC Dashboard; this large number of published results allows users to quickly examine the role of particular biological response components, such as individual genes or cell types, across studies. Most of the currently curated gene signatures have fewer than 50 genes, with a range of 1 to 2,036 (Fig. 3a), while most cell-type frequency or cell activation state signatures have only a single cell type, with a range up to 9 (Fig. 3b). These signatures represent findings from 16 tissues and tissue extracts, including blood, PBMCs, T cells, B cells, monocytes, and NK cells. Nearly 250 entries describe changes over time, more than 75 capture antibody response-associated signatures, and several others come from studies that report effects of age and T cell responses.

The frequency with which cell types and genes are reported in the Dashboard offers insights about key players of the human immune response to vaccination. The most commonly reported genes are STAT1, a key mediator of immune response activated by cytokines and interferons; GBP1, an interferon induced gene involved in innate immunity; IFI44L, a paralog of Interferon Stimulated Protein 44, and SERPING1, a complement cascade protein (Fig. 3c). The most common cell types across pathogens and comparisons are NK cells, CD4+ T cells, and CD8+ T cells (Fig. 3d). We searched the HIPC Dashboard data for genes with vaccination signatures across six or more pathogens and found a set of 36 associated genes across 12 target pathogens (Fig. 3e). Many are interferon stimulated genes, Toll-like receptors, or members of the complement cascade, potentially reflecting a common transcriptional program in response to many different vaccinations.

By design, our current implementation captures vaccination signatures as they are reported in the literature, but it does not include related methodological or statistical information regarding signature discovery (e.g. p-value cut-off) or provide analytical tools that can be applied to the curated signatures. Test statistics are not usually comparable across study designs, and we believe this information may give users a false sense that some signatures are more statistically reliable than others. We instead defer to the judgement of each study's authors and their peer reviewers, and capture signatures as they were reported in each publication. We also caution that bias regarding the number of times particular genes or cell types were investigated might skew relationships in the Dashboard, thus precluding certain types of analyses. As a result, high level analytical tools have not been integrated into the Dashboard, although all of the signatures with full metadata can be downloaded to enable the use of third-party tools. Despite this, we believe the signatures available in the HIPC Dashboard will allow the research community to quickly query the literature and provide valuable comparisons and context for their own experimental results.

The number of genes and cell types captured reflects publications curated through January 2021, but we anticipate the HIPC Dashboard will undergo regular updates to accommodate new findings and additional domains of interest. The current implementation includes gene and cell type response components, as these represent the most commonly published signature types, but it will be valuable to also curate other response components, such as pathways, proteins, and metabolites. Additionally, we recognize that researchers may wish to compare a vaccine response against a particular pathogen to its corresponding disease response; it is easy to see how future iterations could expand the existing vaccine signature framework to capture signatures of infection. Based on our experience, we expanded the data model to include figure numbers or supplementary file annotations within publications as this can greatly simplify quality control during manual curation. We have provided links in the Dashboard to original sources wherever possible. We are also keen on exploring advancements in text-mining and artificial intelligence (AI), to assess how they can assist in automating signature identification and coding. To that end, the immune signatures in the HIPC Dashboard can be used as a data source for training/testing such AI solutions in the future.

In summary, we present the HIPC Dashboard (hipc-dashboard.org) to provide the vaccine research community with easy access to hundreds of published human systems vaccinology signatures. This resource will allow researchers to rapidly compare their own experimental results against existing findings that may otherwise be difficult to locate in the literature. This resource encourages the re-use of published results for advancing our understanding of human vaccine responses and provides a framework that can be extended to capture signatures from other types of immune exposures.

Methods

Manual curation. The initial list of publications to curate into the HIPC Dashboard were derived from a PubMed search of papers matching the terms “Vaccine [AND] Signatures” or “Vaccine [AND] Gene expression”. Publications were further filtered to meet a set of inclusion criteria: (i) study involved human subjects, (ii) provided a comparison of a measurable change or correlation before and after vaccination (or challenge), and (iii) were reported as statistically significant. Signatures were excluded if they were missing directionality, or if they were derived from datasets external to the publication, to avoid redundancy. Two data curators manually collected a standard set of information from each study according to the designed data model (see Fig. 1) and recorded it into a spreadsheet. Each signature was entered by one curator, and subsequently double-checked by the second curator. Table 3 shows a representative portion of the standardized information captured for each signature (12 data fields out of a total of 25).

Assays in the curated publications included gene expression analysis and measured changes in cell-type frequency and cell activation state. Each publication could give rise to any number and type of individual signatures. The signature content is centered on a list of biological response components (genes or cell types) that had a statistically significant change in the assay. These were designated as “response components” to capture different types of entities in a single standardized template column. For example, for gene expression assays, this was often a list of differentially expressed genes.




Fig. 3 Summarization of HIPC Dashboard contents. Signature size distributions showing the number of response components across (a) gene and (b) cell type signatures. (c) Word cloud of the top 50 most frequent gene symbols and (d) top 10 most frequent cell types, where size corresponds to the total number of observations in the Dashboard. (e) Heatmap of recurring genes across vaccines targeting different pathogens. Temporally associated genes in adult whole blood or PBMCs were filtered to those with signatures for six or more pathogens. Color indicates up (red) or down (blue) regulation. Genes with opposing directions in multiple studies were marked ‘trends up’ or ‘trends down’ according to the most common direction (or marked ‘no consensus’ for perfect ties).

For genes, an initial manual curation process was applied to make a first pass at symbol standardization and detect any mistakes in copying. Gene names, symbols and or IDs (which may include HGNC symbols, Entrez IDs, Ensembl IDs etc.) were searched in turn against Panther, (www.pantherdb.org)²², using the “Functional classification viewed in gene list” search, followed if needed by searches against UniProt (www.uniprot.org)²³ and NCBI (www.ncbi.nlm.nih.gov/search) until either a match or updated symbol was found; in the case of no match, the original representation was left unchanged. Any IDs that matched entries that were deprecated or defined as pseudogenes were removed from the curation.

Data standardization. The manually curated data required further steps to match terms encountered to their appropriate ontology representations. A number of the translations described below were orchestrated using an R script, which generated files ready for loading into the HIPC Dashboard.

Gene symbols. Outdated gene symbols and known aliases were translated to their current NCBI representation, which is the HGNC symbol in all but one case. The first pass of conversion used the function `alias2SymbolUsing-NCBI()` from the Bioconductor `limma` package²⁴ with the most recent available gene annotation file. This function returns either an exactly matching official symbol, or if none, the alias with the lowest EntrezID. We followed this by a second R package, `HGNChelper`²⁵, which was able to resolve additional unmatched gene symbols to valid NCBI symbols. Genbank accession numbers were converted to gene symbols where possible using the `org.Hs.egACCNUM2EG` translation table which is part of the Bioconductor `org.Hs.eg.db` package²⁶. Selected symbols still not matching NCBI names were investigated and corrected manually where possible after checking the original publications for context or for errors in transcription. Symbols for which no valid NCBI gene symbol was found, e.g. some pseudogenes, antisense, or uncharacterized genes, are not included in the HIPC Dashboard proper, since a requirement of the Dashboard framework is that all gene symbols must appear in the controlled vocabulary (NCBI/HGNC). However, these symbols are included in downloadable complete gene lists for each signature.

Cell types. Cell types as response components were first curated from the publications as published using a combination of cell type terms and additional descriptive terms, such as protein marker expression. This information was then mapped to a combination of Cell Ontology and Protein Ontology terms, according to a published model¹⁶. Note that cell types can appear in two different contexts, either as response components themselves, or as the cell type isolated for gene expression experiments. In some cases, additional information was provided which could not be mapped to an ontology term¹⁶. This type of information related to a wide variety of cell identification techniques and included the use of additional stains such as viability dyes or tetramer staining. For each set of terms, an entry was created in a lookup table by assigning (1) a parent cell type from the cell ontology, (2) mapping additional protein marker terms to the protein ontology, and then (3) separately retaining as free text descriptors not mapped to an ontology entry, such as tetramer specificity. Thus, the original entry was mapped to up to three descriptor columns, which can be combined as needed for display purposes. For a full, translated cell type, the displayed format is the cell ontology name followed by, if there are additional terms, the “&” symbol, followed by any PRO terms and then any free text.

Vaccines. Vaccine names collected from the literature were manually mapped to the most specific vaccine ontology term available. If a specific vaccine could not be found in VO, new terms were requested. Some examples of terms we included were VO_0004899 (2012–2013 seasonal trivalent inactivated influenza vaccine), VO_0003961 (ChAd63-KH vaccine, *Leishmania donovani*), VO_0004890 (gH1-Qbeta vaccine, novel pandemic-influenza), and VO_0004891 (CN54gp140 + GLA, HIV-1).

Pathogens. The viral, bacterial or protozoan pathogens targeted by each vaccine are represented with terms from the NCBI Taxonomy. For the case of influenza vaccines, a table was created mapping vaccines by year of administration and type (e.g. trivalent or quadrivalent) to their seasonal viral components, unless otherwise indicated in the publication (e.g. for monovalent or specialized vaccines such as Pandemrix). For the few cases where the exact viral strain was not present in the NCBI Taxonomy, the closest more general term in the hierarchy was used. This mapping table was used to substitute in the actual viral pathogen names.

Data availability

Curated signatures are available on the HIPC Dashboard website (<http://hipc-dashboard.org>) and GitHub (see Code Availability for details). Files listing all response components for a signature can be downloaded from within individual observations in the Dashboard. The complete set of signature data can be downloaded from the GitHub repository at <https://github.com/floratos-lab/hipc-dashboard-pipeline>. The data in this paper corresponds to pipeline and data version 1.2.1 in the pipeline GitHub repository²⁷. The GitHub repository contains copies of (1) the original curated data sheets, (2) the response components in individual files, one per signature, (3) the response components in the Broad GMT format⁸, and (4) the actual tab-delimited Dashboard load files, in which the complete signature data is fully denormalized into an easy-to-parse format. Further details about the file formats are available on the GitHub project page. R session information for the Dashboard signature pre-processing pipeline is available on GitHub.

Code availability

Source curated data and mapping files (cell types, vaccine components, the NCBI gene file used, etc.), as well as the R code for the processing pipeline used to create the Dashboard submission files, are available on GitHub at <https://github.com/floratos-lab/hipc-dashboard-pipeline>.

Supported web browsers - The HIPC Dashboard has been tested on recent versions of

- Chrome (Version 93.0.4577.63 (Official Build) (64-bit))
- Firefox (Version 92.0 (64-bit))
- Edge (Version 93.0.961.38 (Official build) (64-bit))

Received: 4 May 2021; Accepted: 14 July 2022;

Published online: 08 November 2022

References

1. Edgar, R., Domrachev, M. & Lash, A. E. Gene Expression Omnibus: NCBI gene expression and hybridization array data repository. *Nucleic Acids Res* **30**, 207–210 (2002).
2. Bhattacharya, S. *et al.* ImmPort, toward repurposing of open access immunological assay data for translational and clinical research. *Sci Data* **5**, 180015 (2018).
3. Bucasas, K. L. *et al.* Early patterns of gene expression correlate with the humoral immune response to influenza vaccination in humans. *The Journal of infectious diseases* **203**, 921–929 (2011).
4. Shah, N. *et al.* A crowdsourcing approach for reusing and meta-analyzing gene expression data. *Nat. Biotechnol.* **34**, 803–806 (2016).
5. Chen, E. Y. *et al.* Enrichr: interactive and collaborative HTML5 gene list enrichment analysis tool. *BMC Bioinformatics* **14**, 128 (2013).
6. Kuleshov, M. V. *et al.* Enrichr: a comprehensive gene set enrichment analysis web server 2016 update. *Nucleic Acids Res.* **44**, W90–97 (2016).
7. Pilarczyk, M. *et al.* Connecting omics signatures of diseases, drugs, and mechanisms of actions with iLINCS. 826271 <https://doi.org/10.1101/826271> (2020).
8. Subramanian, A. *et al.* Gene set enrichment analysis: A knowledge-based approach for interpreting genome-wide expression profiles. *PNAS* **102**, 15545–15550 (2005).
9. Liberzon, A. *et al.* The Molecular Signatures Database (MSigDB) hallmark gene set collection. *Cell Syst* **1**, 417–425 (2015).
10. Berke, K. *et al.* VaximmutorDB: A Web-Based Vaccine Immune Factor Database and Its Application for Understanding Vaccine-Induced Immune Mechanisms. *Frontiers in Immunology* **12**, 645 (2021).
11. Aksoy, B. A. *et al.* CTD2 Dashboard: a searchable web interface to connect validated results from the Cancer Target Discovery and Development Network. *Database (Oxford)* **2017** (2017).
12. Yates, B. *et al.* Genenames.org: the HGNC and VGNC resources in 2017. *Nucleic Acids Res.* **45**, D619–D625 (2017).
13. Bard, J., Rhee, S. Y. & Ashburner, M. An ontology for cell types. *Genome Biology* **6**, R21 (2005).
14. Diehl, A. D. *et al.* The Cell Ontology 2016: enhanced content, modularization, and ontology interoperability. *J Biomed Semant* **7**, 44 (2016).
15. Natale, D. A. *et al.* Protein Ontology (PRO): enhancing and scaling up the representation of protein entities. *Nucleic Acids Res.* **45**, D339–D346 (2017).
16. Overton, J. A. *et al.* Reporting and connecting cell type names and gating definitions through ontologies. *BMC Bioinformatics* **20**, 182 (2019).
17. Vita, R. *et al.* A structured model for immune exposures. *Database (Oxford)* **2020** (2020).
18. He, Y. *et al.* VO: Vaccine Ontology. *Nature Precedings* 1–1 <https://doi.org/10.1038/npre.2009.3552.1> (2009).
19. Sayers, E. W. *et al.* GenBank. *Nucleic Acids Res* **47**, D94–D99 (2019).
20. Schoch, C. L. *et al.* NCBI Taxonomy: a comprehensive update on curation, resources and tools. *Database (Oxford)* **2020** (2020).
21. Vita, R. *et al.* The Immune Epitope Database (IEDB): 2018 update. *Nucleic Acids Res.* **47**, D339–D343 (2019).
22. Mi, H., Poudel, S., Muruganujan, A., Casagrande, J. T. & Thomas, P. D. PANTHER version 10: expanded protein families and functions, and analysis tools. *Nucleic Acids Res* **44**, D336–D342 (2016).
23. The UniProt Consortium. UniProt: a worldwide hub of protein knowledge. *Nucleic Acids Research* **47**, D506–D515 (2019).
24. Ritchie, M. E. *et al.* limma powers differential expression analyses for RNA-sequencing and microarray studies. *Nucleic Acids Research* **43**, e47 (2015).
25. Oh, S. *et al.* HGNCHELPER: identification and correction of invalid gene symbols for human and mouse. *F1000Res* **9**, 1493 (2020).
26. Carlson, M. org.Hs.eg.db: Genome wide annotation for Human. R package version 3.14.0. (2021).
27. Smith, K. & Ji, Z. floratos-lab/hipc-dashboard-pipeline: HIPC Dashboard pipeline v1.2.1. *Zenodo* <https://doi.org/10.5281/zenodo.5656986> (2021).

Acknowledgements

This research was performed as a project of the Human Immunology Project Consortium (HIPC) and supported by the National Institute of Allergy and Infectious Diseases. This work was supported in part by NIH grants U19AI128949, U19AI118608, U19AI118626, and U19AI089992. This work was supported in part by the Canadian Institutes of Health Research [funding reference number FDN-154287].

Author contributions

Conceptualization: A.F., S.H.K., H.I.P.C.; Methodology: K.C.S., D.C., B.K.D., Z.J., R.V., B.P., R.E.W.H., A.F., S.H.K.; Software: K.C.S., Z.J., A.F.; Investigation: K.C.S., D.C., B.K.D., E.v.d.L., J.Y.J.W., E.T., A.A.; Writing - Original Draft: K.C.S., D.C., B.K.D., A.F., S.H.K.; Writing - Review and Editing: all authors.

Competing interests

S.H.K. receives consulting fees from Northrop Grumman and Peraton. The remaining authors declare no competing interests.

Additional information

Supplementary information The online version contains supplementary material available at <https://doi.org/10.1038/s41597-022-01558-1>.

Correspondence and requests for materials should be addressed to A.F. or S.H.K.

Reprints and permissions information is available at www.nature.com/reprints.

Publisher's note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.



Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons license, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons license and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this license, visit <http://creativecommons.org/licenses/by/4.0/>.

© The Author(s) 2022

The Human Immunology Project Consortium (HIPC)

Bjoern Peters^{4,5}, Steven H. Kleinstein^{2,7,8,10}✉

A full list of members and their affiliations appears in the Supplementary Information.