



OPEN

Optimizing nutrient use efficiency, productivity, energetics, and economics of red cabbage following mineral fertilization and biopriming with compatible rhizosphere microbes

Deepranjan Sarkar¹, Ardith Sankar², O. Siva Devika¹, Sonam Singh¹, Shikha³, Manoj Parihar⁴, Amitava Rakshit¹✉, R. Z. Sayyed⁵, Abdul Gafur⁶, Mohammad Javed Ansari⁷, Subhan Danish^{8,10}✉, Shah Fahad⁹ & Rahul Datta¹⁰

Conventional agricultural practices and rising energy crisis create a question about the sustainability of the present-day food production system. Nutrient exhaustive crops can have a severe impact on native soil fertility by causing nutrient mining. In this backdrop, we conducted a comprehensive assessment of bio-priming intervention in red cabbage production considering nutrient uptake, the annual change in soil fertility, nutrient use efficiency, energy budgeting, and economic benefits for its sustainable intensification, among resource-poor farmers of Middle Gangetic Plains. The compatible microbial agents used in the study include *Trichoderma harzianum*, *Pseudomonas fluorescens*, and *Bacillus subtilis*. Field assays (2016–2017 and 2017–2018) of the present study revealed supplementing 75% of recommended NPK fertilizer with dual inoculation of *T. harzianum* and *P. fluorescens* increased macronutrient uptake (N, P, and K), root length, heading percentage, head diameter, head weight, and the total weight of red cabbage along with a positive annual change in soil organic carbon. Maximum positive annual change in available N and available P was recorded under 75% RDF + *P. fluorescens* + *B. subtilis* and 75% RDF + *T. harzianum* + *B. subtilis*, respectively. Bio-primed plants were also higher in terms of growth and nutrient use efficiency (agronomic efficiency, physiological efficiency, apparent recovery efficiency, partial factor productivity). Energy output (26,370 and 26,630 MJ ha⁻¹), energy balance (13,643 and 13,903 MJ ha⁻¹), maximum gross return (US \$ 16,030 and 13,877 ha⁻¹), and net return (US \$ 15,966 and 13,813 ha⁻¹) were considerably higher in *T. harzianum*, and *P. fluorescens* treated plants. The results suggest the significance of the bio-priming approach under existing integrated nutrient management strategies and the role of dual inoculations in producing synergistic effects on plant growth and maintaining the soil, food, and energy nexus.

¹Department of Soil Science and Agricultural Chemistry, Institute of Agricultural Sciences, Banaras Hindu University, Uttar Pradesh, Varanasi 221005, India. ²Department of Agronomy, Institute of Agricultural Sciences, Banaras Hindu University, Uttar Pradesh, Varanasi 221005, India. ³Krishi Vigyan Kendra, Ranichauri, Veer Chandra Singh Garhwali Uttarakhand University of Horticulture and Forestry, Tehri Garhwal 249199, Uttarakhand, India. ⁴Crop Production Division, ICAR-Vivekananda Parvatiya Krishi Anusandhan Sansthan, Almora 263601, Uttarakhand, India. ⁵Department of Microbiology, PSGVP Mandal's, Arts, Science & Commerce College, 425409, Shahada, Maharashtra, India. ⁶Sinarmas Forestry Corporate Research and Development, Perawang 28772, Indonesia. ⁷Department of Botany, Hindu College Moradabad (Mahatma Jyotiba Phule Rohilkhand University Bareilly), Moradabad, Uttar Pradesh 244001, India. ⁸Department of Soil Science, Faculty of Agricultural Sciences and Technology, Bahauddin Zakariya University, Multan 60800, Pakistan. ⁹Department of Agronomy, The University of Haripur, Haripur 22620, Pakistan. ¹⁰Department of Geology and Pedology, Faculty of Forestry and Wood Technology, Mendel University in Brno, Zemedelska1, 61300 Brno, Czech Republic. ✉email: amitavar@bhu.ac.in; sd96850@gmail.com

India has attained ‘self-sufficiency’ in food grain production, but some factors like declining nutrient use efficiency, low crop response ratio, negative soil nutrient balance, etc., have become major concerns in Indian agriculture. The accomplishment of the Green Revolution has left some drastic effects on our motherland soil. As we fulfilled our target of increased food grain production, there are indications of declining partial and total factor productivity, and the sole cause has been attributed to diminishing native soil fertility¹. Therefore, the dwindling natural resource base has raised sustainability concerns and integrated resource management in the present agricultural system. Moreover, climate change has forced us to adopt crop management practices more scientifically to sustainable land use and land cover².

Beneficial microorganisms play an important role in achieving agro-environmental sustainability. However, the abundance and robustness of microbes are dependent upon smart delivery systems³. Bio-priming is one possible method to apply plant growth-promoting microorganisms in the soil–plant system and improve the nutrient use efficiency of agroecosystems^{4,5}. Primers’ selection is an essential step in biological seed enhancements as priming agents’ growth-promoting abilities are highly specific to certain species, cultivars, or genotypes of crops⁶. A microbial consortium treatment can also be used for getting better results after checking the compatibility of the microbes⁷. This will promote synergistic interactions among the microbial population⁸. The literature of current decades reveals that beneficial microorganisms and their interactions with host plants harbor broad prospects in sustainable agriculture’s persistence. Their influence in the rhizosphere ecosystem compels researchers to investigate their ecology, diversity, and activity, contributing most to soil health and plant fitness. The rhizosphere is chosen because it is the hub of a complex web of interactions regulated by the system’s enormous energy flux, releasing about 20–50% of plant photosynthates from the roots^{9,10}. Thus, strong documentation is required on how species richness affects the ecological functions of agroecosystems. Understanding the microbial mechanisms influencing plant productivity is key to augment plant growth and boost the processes within the soil system. In general, the direct growth-promoting mechanisms used by them can be listed as (i) nitrogen (N) fixation; (ii) nutrient solubilization (phosphorus (P), potassium (K), etc.); (iii) production of phytohormones, e.g., auxins (indole-3-acetic acid, indole-3-butyric acid), cytokinins, gibberellins, abscisic acid, and ethylene; (iv) iron sequestration by siderophore production; (v) regulation of plant hormones, e.g., 1-aminocyclopropane-1-carboxylate (ACC) deaminase decreases ethylene levels; and the indirect ones include (i) induction of systemic resistance against plant pathogens through antagonism with the production of inhibitory compounds or substances; (ii) suppression of phytopathogens with the synthesis of hydrolytic enzymes (glucanases, chitinases, proteases, and lipases), siderophores, antibiotics, and cyanide^{11–14}.

The application of fertilizers for crop production is increasing rapidly. However, the average N recovery efficiency and P recovery efficiency are less than 50% and 20%, respectively¹⁵. Some recent studies showed improved nutrient use efficiency due to microbial inoculation of plants. Soils treated with 50% organic and inorganic N fertilizers and the application of *Paraburkholderia* sp. augmented the growth of kikuyu grass and N use efficiency compared to 100% N fertilizer treatment¹⁶. Pereira et al.¹⁷ reported improved N use efficiency and P use efficiency of maize plants treated with *Cupriavidus necator* and *Pseudomonas fluorescens*. However, these experiments were performed under pot conditions.

The modern agricultural system depends on energy inputs, viz., fossil fuels, fertilizers, pesticides, electricity, etc. that affect our ecosystem’s health and increase greenhouse gas emissions¹⁸. The extent of energy consumption is evident from the fact that the production of nitrogen (N), phosphorus (P), and potassium (K) fertilizers require approximately 60.6, 11.1, and 6.7 MJ kg⁻¹ energy, respectively^{19,20}. However, very few studies have evaluated the energetics of crop production involving microbial application, mainly based on controlled conditions. Mihov and Tringovska²¹ evaluated the energy use efficiency of greenhouse tomato production. Pal and Singh²² examined the relationship of energy inputs and crop yield in greenhouse okra production.

With growing agro-environmental challenges, we need to look for new, nutritive, and versatile crops to make agriculture sustainable and profitable. Red cabbage is an excellent source of minerals, vitamins, antioxidants, anthocyanin, glucosinolates, and phenolic compounds^{23,24}. Therefore, our diet is often chosen as salads, micro-greens, leafy vegetables, and health-improving supplements. In addition, supermarkets will provide assured economic returns to farmers. Hence, the present study’s primary objective was to investigate the effect of bio-priming and mineral fertilization on red cabbage growth, nutrient uptake, the annual change in soil fertility, nutrient use efficiency, energy usage, and economic returns under field conditions.

Results and discussion

Macronutrient uptake. Total uptake of N, P, and K by red cabbage was partitioned into head and stalk. The head’s highest N uptake was registered in plots receiving 75% RDF + *T. harzianum* + *P. fluorescens* (T₆) during both the years (Table S1). It showed 29% (1st year) and 24% (2nd year) increments over 100% RDF (T₂). Similarly, a significant increase in total uptake of N was also observed in T₆ treatment. The total N uptake varied from 21.53 to 70.21 kg ha⁻¹ in the first year and 19.56–69.96 kg ha⁻¹ in the second year. Compared with single-priming and consortium treatments, N total uptake increased by 12–35% with the co-application of *T. harzianum* and *P. fluorescens*. The uptake of the other two dual bio-priming treatments, viz., T₇ (75% RDF + *P. fluorescens* + *B. subtilis*) and T₈ (75% RDF + *T. harzianum* + *B. subtilis*) were at par with T₂ (100% RDF). Phosphorus uptake by red cabbage was also influenced by 75% RDF + *T. harzianum* + *P. fluorescens* (T₆), but it was at par with 75% RDF + *P. fluorescens* + *B. subtilis* (T₇). In the head, P uptake ranged from 1.28 to 7.36 kg ha⁻¹ in the first year and 0.80–7.52 kg ha⁻¹ in the second year (Table S2). During the second year, the head uptake of plants under 75% RDF + *P. fluorescens* (T₄) was higher than 75% RDF + *T. harzianum* + *B. subtilis* (T₈). The application of fertilizers and bio-priming agents did not significantly affect the P uptake by stalk during the study years. The total P uptake varied from 3.69 to 10.99 kg ha⁻¹ in the first year and 3.44–11.29 kg ha⁻¹ in the second year. Sole application of *T. harzianum* and *B. subtilis* and a triple consortium of *T. harzianum*, *P. fluorescens*, and *B. subtilis* showed statistically




Figure 1. Annual change in organic carbon (A), available N (B), available P (C), and available K (D) as affected by bio-priming and mineral fertilization. Error bars indicate mean \pm SE ($n=3$). Treatments: T₁: Absolute control N:P₂O₅:K₂O @ 0:0:0 kg ha⁻¹; T₂: RDF of N:P₂O₅:K₂O @ 120:60:60 kg ha⁻¹; T₃: 75% RDF + *Trichoderma harzianum*; T₄: 75% RDF + *Pseudomonas fluorescens*; T₅: 75% RDF + *Bacillus subtilis*; T₆: 75% RDF + *T. harzianum* + *P. fluorescens*; T₇: 75% RDF + *P. fluorescens* + *B. subtilis*; T₈: 75% RDF + *T. harzianum* + *B. subtilis*; T₉: 75% RDF + *T. harzianum* + *P. fluorescens* + *B. subtilis*.

similar P uptake. Except for control, the uptake was found to increase in all the treatments during the second year over 1st year. In the case of K, a significant increase in uptake (head and total) was recorded in T₆ treatment (Table S3). The magnitude of the increase due to this treatment over 100% RDF (T₂) was 31% (head) and 21% (total). However, applying 100% chemical fertilizers and combined use of chemical fertilizers and bio-agents did not significantly differ in the uptake of macronutrients by the stalk. In general, the K uptake varied from 8.25 to 57.35 kg ha⁻¹ for the head and 16.64–22.15 kg ha⁻¹ for the stalk. Stalk K uptake was highest (22.15 kg ha⁻¹) in the first year with 75% RDF + *B. subtilis* (T₅). The total K uptake ranged from 29.03 to 78.20 kg ha⁻¹ in the first year and 26.08–78.51 kg ha⁻¹ in the second year. During the second year, the total K uptake of plants under 100% RDF was par with single- and triple-priming treatments. Among single- and triple-priming treatments, the highest total K uptake was observed with the application of 75% RDF + *P. fluorescens* (T₄) during both years. The lowest N, P, and K uptake were recorded from the control (T₁) plots. On average, red cabbage removed macronutrients in the order of K (64.32 kg ha⁻¹) > N (55.05 kg ha⁻¹) > P (8.91 kg ha⁻¹). The higher nutrient uptake in the integrated application of chemical fertilizers and bio-agents is explained by developing proliferous root systems in bio-primed plants and increased microbial activity in the soil, which helped mineralize nutrients maintaining the soil solution greater assimilation in plants. Several workers^{25–31} reported increased nutrient uptake in crops due to the combined application of chemical fertilizers (reduced level) and biofertilizers.

Annual change in organic carbon (OC) and available N, P, and K. A positive annual change in OC (Fig. 1) was noted for all the treatments (T₂–T₉) except in control plots (T₁) which showed a negative change of -0.22 g kg⁻¹ year⁻¹. The highest positive change was documented under 75% RDF + *T. harzianum* + *P. fluorescens* (T₆, 0.59 g kg⁻¹ year⁻¹) followed by 75% RDF + *P. fluorescens* + *B. subtilis* (T₇, 0.49 g kg⁻¹ year⁻¹), and the lowest was recorded under 100% chemical fertilization (T₂) being 0.17 g kg⁻¹ year⁻¹ which was at par with 75% RDF + *T. harzianum* + *P. fluorescens* + *B. subtilis* (T₉, 0.20 g kg⁻¹ year⁻¹). In the case of annual change of available N, P, and K, similar trends were noticed as OC (Fig. 1). However, the highest and lowest changes varied. Application of 75% RDF + *P. fluorescens* + *B. subtilis* (T₇) and 75% RDF + *T. harzianum* + *B. subtilis* (T₈) demonstrated (Fig. 1) higher positive changes in available N (11.06 and 10.06 kg ha⁻¹ year⁻¹, respectively), while 100% RDF (T₂) showed the lowest change (4.47 kg ha⁻¹ year⁻¹). The control plots (T₁) showed a significant negative annual change in available N (-13.46 kg ha⁻¹ year⁻¹) after crop harvest. The T₈ treatment showed the maximum positive annual change (6.70 kg ha⁻¹ year⁻¹) followed by T₇ (6.27 kg ha⁻¹ year⁻¹) for available P (Fig. 1). However, the least positive change in available P was evident in T₂ (3.10 kg ha⁻¹ year⁻¹), which was at par with the treatment T₉ (3.48 kg ha⁻¹ year⁻¹). A negative change of -1.27 kg ha⁻¹ year⁻¹ in available P was observed in T₁ plots. There were no significant positive annual changes observed between the treatments for available K (Fig. 1). However, the T₈ treatment presented the highest positive annual change value of 10.04 kg ha⁻¹ year⁻¹, followed by T₇ (8.92 kg ha⁻¹ year⁻¹); the rest of the treatments (T₂, T₃, T₄, T₅, T₆, and T₉) documented at par values of the same. The T₁ plots recorded a significant negative annual change in available K (-11.43 kg ha⁻¹ year⁻¹).

Treatments	N use efficiency							
	Agronomic efficiency (kg kg ⁻¹)		Physiological efficiency (kg kg ⁻¹)		Apparent recovery efficiency (%)		Partial factor productivity (kg kg ⁻¹)	
	2016–2017	2017–2018	2016–20	2017–2018	2016–2017	2017–2018	2016–2017	2017–2018
T ₁ : Absolute control N:P ₂ O ₅ :K ₂ O @ 0:0:0 kg ha ⁻¹	–	–	–	–	–	–	–	–
T ₂ : RDF of N:P ₂ O ₅ :K ₂ O @ 120:60:60 kg ha ⁻¹	9.48 ^c	11.07 ^c	28.59	33.21	32.73 ^b	33.21 ^d	13.97 ^c	14.05 ^c
T ₃ : 75% RDF + <i>Trichoderma harzianum</i>	12.06 ^{bc}	15.02 ^{abc}	32.33	34.57	37.38 ^b	43.39 ^{bc}	18.06 ^{bc}	18.99 ^{ab}
T ₄ : 75% RDF + <i>Pseudomonas fluorescens</i>	13.09 ^{abc}	15.80 ^{ab}	31.98	36.53	41.09 ^b	43.49 ^{bc}	19.08 ^{ab}	19.77 ^{ab}
T ₅ : 75% RDF + <i>Bacillus subtilis</i>	11.45 ^{bc}	14.55 ^{bc}	33.74	37.49	34.03 ^b	38.83 ^{cd}	17.44 ^{bc}	18.52 ^b
T ₆ : 75% RDF + <i>T. harzianum</i> + <i>P. fluorescens</i>	17.44 ^a	19.38 ^a	32.02	34.48	54.10 ^a	55.99 ^a	23.44 ^a	23.34 ^a
T ₇ : 75% RDF + <i>P. fluorescens</i> + <i>B. subtilis</i>	15.44 ^{ab}	17.52 ^{ab}	34.06	39.27	40.10 ^b	44.57 ^{bc}	21.44 ^{ab}	21.49 ^{ab}
T ₈ : 75% RDF + <i>T. harzianum</i> + <i>B. subtilis</i>	13.69 ^{abc}	16.70 ^{ab}	30.96	34.75	36.76 ^b	48.01 ^b	19.68 ^{ab}	20.67 ^{ab}
T ₉ : 75% RDF + <i>T. harzianum</i> + <i>P. fluorescens</i> + <i>B. subtilis</i>	10.96 ^{bc}	13.97 ^{bc}	32.58	35.89	33.83 ^b	38.33 ^{cd}	16.95 ^{bc}	17.94 ^{bc}

Table 1. Nitrogen use efficiency of red cabbage as influenced by bio-priming and fertilisation. Different letters indicate significant differences at $P \leq 0.05$ among the treatments as per DMRT.

Conclusively, due to no supplement of fertilizer nutrients and/or biofertilizers, the control plots exhibited a negative annual change in all the studied cases demonstrating the loss/mining of nutrients. However, where both are supplied, a positive change is noticed, indicating annual enrichment of the respected soil attributes over control. In this study, T₈ and T₇ treatments performed better in enhancing available N, P, and K except in OC where T₆ treatment presented a significant positive result over these two, and the rest of the treatments remained at par with positive change. Furthermore, Kaur and Reddy³² reported improved soil fertility using mineral fertilizer and biofertilizer (*Pseudomonas plecoglossicida* and *Pantoea cyripedii*). Similarly, the integrated application of NPK fertilizers and organic amendments (green manure + *Pseudomonas putida* + *Azotobacter chroococcum*) recorded a higher OC value and available N and K over only NPK fertilizer application³³.

Nitrogen use efficiency. Optimization of fertilizer for achieving higher input use efficiency without hindering the economic yield is a crucial issue in agriculture. The extent of crop utilization to applied N was analyzed through different parameters of N use efficiency. Agronomic efficiency (AE_N) varied from 9.48 to 19.38 kg of head kg⁻¹ of N applied (Table 1). During the study period of the field experiment, the mean value of AE_N increased from 12.95 kg of head kg⁻¹ of N applied in 2016–2017 to 15.50 kg of head kg⁻¹ of N applied in 2017–2018. The application of 75% RDF + *T. harzianum* + *P. fluorescens* (T₆) increased AEN by 84% and 75% over 100% RDF (T₂) during the first and second years. Compared to T₆, the application of 75% RDF + *T. harzianum* + *P. fluorescens* + *B. subtilis* (T₉) significantly reduced the AE_N by 59% and 39% in the first and second years respectively. Application of triple consortium also recorded low AE_N in comparison to single-species bio-priming and dual microbial consortium treatments. Seedling inoculation with biofertilizers along with 75% RDF + vermicompost demonstrated similar results in cabbage³⁴.

Physiological efficiency (PE_N) reflects better accumulation and conversion of N from source to sink. The highest PE_N (34.06 and 39.27 kg kg⁻¹) was observed with the application of 75% RDF + *P. fluorescens* + *B. subtilis* (T₇) during both the years of study (Table 1). Low PE_N of crops was recorded under 75% RDF + *T. harzianum* + *P. fluorescens* (T₆). Plants under 75% RDF + *B. subtilis* (T₅) registered 4% higher PE_N compared to that under 75% RDF + *T. harzianum* + *P. fluorescens* + *B. subtilis* (T₉). Among the single-species bio-priming agents, *B. subtilis* resulted in maximum PE_N (33.74 and 37.49 kg kg⁻¹) during both years. Individual application of *T. harzianum* and co-application of *T. harzianum* and *P. fluorescens* showed an equivalent effect on PE_N.

Nitrogen use efficiency in apparent recovery efficiency (ARE_N) varied from 32.73 to 55.99% (Table 1). The highest N use efficiency of 54.10% and 55.99% in 2016–2017 and 2017–2018, respectively, was observed under 75% RDF + *T. harzianum* + *P. fluorescens* (T₆). Among the bio-priming treatments, the lowest ARE_N was obtained under triple consortium treatment (T₉). In the first year, ARE_N in *P. fluorescens* bio-primed plants was greater than other individual bio-priming agents. However, in the second year, single bio-priming of *T. harzianum* and *P. fluorescens* recorded similar N use efficiency. In addition, improved ARE_N was reported in lettuce plants treated with *Trichoderma*-based biostimulants³⁵.

Regarding partial factor productivity (PFP_N), the results showed similar trends to that of ARE_N. The response ranged between 13.97 and 23.44 kg of head kg⁻¹ of N applied (Table 1). Application of 75% RDF + *T. harzianum* + *P. fluorescens* (T₆) and 75% RDF + *T. harzianum* + *P. fluorescens* + *B. subtilis* (T₉) registered highest and

Treatments	P use efficiency							
	Agronomic efficiency (kg kg ⁻¹)		Physiological efficiency (kg kg ⁻¹)		Apparent recovery efficiency (%)		Partial factor productivity (kg kg ⁻¹)	
	2016–2017	2017–2018	2016–2017	2017–2018	2016–2017	2017–2018	2016–2017	2017–2018
T ₁ : Absolute control N:P ₂ O ₅ :K ₂ O @ 0:0:0 kg ha ⁻¹	–	–	–	–	–	–	–	–
T ₂ : RDF of N:P ₂ O ₅ :K ₂ O @ 120:60:60 kg ha ⁻¹	18.96 ^c	22.15 ^c	227.25	233.68 ^{ab}	8.18 ^c	9.50 ^c	27.94 ^c	28.10 ^c
T ₃ : 75% RDF + <i>Trichoderma harzianum</i>	24.13 ^{bc}	30.04 ^{abc}	231.84	241.04 ^a	10.68 ^{bc}	12.46 ^{bc}	36.11 ^{bc}	37.97 ^{ab}
T ₄ : 75% RDF + <i>Pseudomonas fluorescens</i>	26.18 ^{abc}	31.60 ^{ab}	206.31	207.53 ^b	12.89 ^{ab}	15.25 ^{ab}	38.16 ^{ab}	39.54 ^{ab}
T ₅ : 75% RDF + <i>Bacillus subtilis</i>	22.91 ^{bc}	29.10 ^{bc}	201.95	241.60 ^a	11.30 ^{bc}	12.22 ^{bc}	34.89 ^{bc}	37.04 ^b
T ₆ : 75% RDF + <i>T. harzianum</i> + <i>P. fluorescens</i>	34.89 ^a	38.75 ^a	213.14	221.02 ^{ab}	16.18 ^a	17.45 ^a	46.87 ^a	46.69 ^a
T ₇ : 75% RDF + <i>P. fluorescens</i> + <i>B. subtilis</i>	30.89 ^{ab}	35.05 ^{ab}	190.59	208.97 ^b	16.23 ^a	16.95 ^a	42.87 ^{ab}	42.98 ^{ab}
T ₈ : 75% RDF + <i>T. harzianum</i> + <i>B. subtilis</i>	27.38 ^{abc}	33.40 ^{ab}	205.72	242.77 ^a	13.23 ^{ab}	13.84 ^{ab}	39.36 ^{ab}	41.33 ^{ab}
T ₉ : 75% RDF + <i>T. harzianum</i> + <i>P. fluorescens</i> + <i>B. subtilis</i>	21.93 ^{bc}	27.94 ^{bc}	215.89	242.77 ^a	10.14 ^{bc}	11.64 ^{bc}	33.91 ^{bc}	35.88 ^{bc}

Table 2. Phosphorus use efficiency of red cabbage as influenced by bio-priming and fertilisation. Different letters indicate significant differences at $P \leq 0.05$ among the treatments as per DMRT.

lowest N use efficiency, respectively. About 68% increase in PFP_N was found in T₆ compared to 100% RDF (T₂). The effect of *P. fluorescens* on PFP_N was higher than other single-species bio-priming agents. The results revealed that dual consortium treatments were more competent in converting the applied N into marketable yield. Our result was consistent with the observation of Chatterjee et al.³⁴.

Phosphorus use efficiency. Agronomic efficiency (AE_p) was maximum under 75% RDF + *T. harzianum* + *P. fluorescens* (T₆) which registered an increment in P use efficiency from 34.89 kg of head kg⁻¹ of P applied in 2016–2017 to 38.75 kg of head kg⁻¹ of P applied in 2017–2018 (Table 2). Application of triple consortium was found to reduce the AE_p in a significant manner. Results showed the following trend: 75% RDF + *T. harzianum* + *P. fluorescens* (T₆) > 75% RDF + *P. fluorescens* + *B. subtilis* (T₇) > 75% RDF + *T. harzianum* + *B. subtilis* (T₈) > 75% RDF + *P. fluorescens* (T₄) > 75% RDF + *T. harzianum* (T₃) > 75% RDF + *B. subtilis* (T₅) > 75% RDF + *T. harzianum* + *P. fluorescens* + *B. subtilis* (T₉) > 100% RDF (T₂). Sole application of *P. fluorescens* along with 75% RDF (T₆) increased the AE_p by 38% and 43% in the first and second year, respectively, compared with 100% RDF (T₂).

Phosphorus use efficiency in physiological efficiency (PE_p) varied from 227.25 to 242.77 kg of head kg⁻¹ of P applied (Table 2). In the first year, application of 75% RDF + *T. harzianum* (T₃) registered maximum PE_p (231.84 kg kg⁻¹) while application of 75% RDF + *T. harzianum* + *B. subtilis* (T₈) and 75% RDF + *T. harzianum* + *P. fluorescens* + *B. subtilis* (T₉) registered maximum PE_p (242.77 kg kg⁻¹) during the second year. Similar to PE_N, PE_p lowered down with application of 75% RDF + *T. harzianum* + *P. fluorescens* (T₆). In the second year, individual application of *P. fluorescens* showed lower PE_p than *B. subtilis* and *T. harzianum*. Yaseen and Malhi³⁶ reported a similar range of PEP in wheat, and they further claimed that P decreased P use efficiency significantly.

The effect of fertilization and bio-priming on apparent recovery efficiency (ARE_p) is presented in Table 2. The ARE_p was found to be maximum (16.23%) under 75% RDF + *P. fluorescens* + *B. subtilis* (T₇) during the first year, but during the second year, maximum (17.45%) ARE_p was noted under 75% RDF + *T. harzianum* + *P. fluorescens* (T₆). Among the three bio-agents, P use efficiency was in *P. fluorescens* > *B. subtilis* > *T. harzianum*. The present study results revealed that bio-priming with *P. fluorescens* demonstrated the highest increment (18%) in ARE_p compared to the first year. Application of triple consortium and single bio-priming with *T. harzianum* resulted in a similar ARE_p. Increased P use efficiency due to co-inoculation of rhizospheric bacterial (endophytic) agents was reported by Emami et al.³⁷.

Partial factor productivity (PFP_p) ranged from 27.94 to 46.87 kg of head kg⁻¹ of P applied with an average value of 37.51 kg of head kg⁻¹ of P applied during the first year and 38.69 kg of head kg⁻¹ of P applied during the second year (Table 2). Among the bio-priming treatments, the application of 75% RDF + *T. harzianum* + *P. fluorescens* (T₆) and 75% RDF + *T. harzianum* + *P. fluorescens* + *B. subtilis* (T₉) recorded the highest and lowest PFP_p, respectively. Compared with 100% RDF (T₂), T₆ increased the P use efficiency by 67%. *P. fluorescens* exhibited the highest P use efficiency regarding individual bio-priming agents, followed by *T. harzianum* and *B. subtilis*. However, compared with the first year, bio-priming with *B. subtilis* showed a 6% increment in PFP_p during the second year. The results of PFP_p followed the order: dual-species bio-priming > single-species bio-priming > triple-species bio-priming.

Treatments	K use efficiency							
	Agronomic efficiency (kg kg ⁻¹)		Physiological efficiency (kg kg ⁻¹)		Apparent recovery efficiency (%)		Partial factor productivity (kg kg ⁻¹)	
	2016–2017	2017–2018	2016–2017	2017–2018	2016–2017	2017–2018	2016–2017	2017–2018
T ₁ : Absolute control N:P ₂ O ₅ :K ₂ O @ 0:0:0 kg ha ⁻¹	–	–	–	–	–	–	–	–
T ₂ : RDF of N:P ₂ O ₅ :K ₂ O @ 120:60:60 kg ha ⁻¹	18.96 ^c	22.15 ^c	32.08	34.33	59.13 ^d	64.27 ^c	27.94 ^c	28.10 ^c
T ₃ : 75% RDF + <i>Trichoderma</i> <i>harzianum</i>	24.13 ^{bc}	30.04 ^{abc}	32.31	33.93	74.45 ^{cd}	88.44 ^{abc}	36.11 ^{bc}	37.97 ^{ab}
T ₄ : 75% RDF + <i>Pseu-</i> <i>domonas fluorescens</i>	26.18 ^{abc}	31.60 ^{ab}	31.25	33.49	84.02 ^{abcd}	94.83 ^{ab}	38.16 ^{ab}	39.54 ^{ab}
T ₅ : 75% RDF + <i>Bacillus</i> <i>subtilis</i>	22.91 ^{bc}	29.10 ^{bc}	29.47	33.29	77.54 ^{bcd}	87.75 ^{abc}	34.89 ^{bc}	37.04 ^b
T ₆ : 75% RDF + <i>T.</i> <i>harzianum</i> + <i>P.</i> <i>fluorescens</i>	34.89 ^a	38.75 ^a	31.68	33.67	109.26 ^a	116.49 ^a	46.87 ^a	46.69 ^a
T ₇ : 75% RDF + <i>P.</i> <i>fluorescens</i> + <i>B.</i> <i>subtilis</i>	30.89 ^{ab}	35.05 ^{ab}	29.18	31.32	105.92 ^{ab}	112.53 ^{ab}	42.87 ^{ab}	42.98 ^{ab}
T ₈ : 75% RDF + <i>T.</i> <i>harzianum</i> + <i>B.</i> <i>subtilis</i>	27.38 ^{abc}	33.40 ^{ab}	29.35	32.09	93.83 ^{abc}	104.45 ^{ab}	39.36 ^{ab}	41.33 ^{ab}
T ₉ : 75% RDF + <i>T.</i> <i>harzianum</i> + <i>P. fluo-</i> <i>rescens</i> + <i>B. subtilis</i>	21.93 ^{bc}	27.94 ^{bc}	30.60	32.69	71.51 ^{cd}	84.97 ^{bc}	33.91 ^{bc}	35.88 ^{bc}

Table 3. Potassium use efficiency of red cabbage as influenced by bio-priming and fertilisation. Different letters indicate significant differences at $P \leq 0.05$ among the treatments as per DMRT.

Potassium use efficiency. The results of K use efficiency in the form of agronomic efficiency (AE_K), and partial factor productivity (PPF_K) was similar to that of P use efficiency (AE_P and PPF_P) because the amount of nutrient applied for P and K was the same, i.e., 60 kg as RDF. The maximum and minimum K use efficiency (AE_K and PPF_K) were obtained from 75% RDF + *T. harzianum* + *P. fluorescens* (T₆) and 100% RDF (T₂), respectively (Table 3). The performance of the triple consortium was lowest among the bio-primed treatments. Application of *P. fluorescens* resulted in higher AE_K in terms of single-species bio-priming.

Physiological efficiency (PE_K) varied from 29.18 to 34.33 kg of head kg⁻¹ of K applied (Table 3). The highest PE_K of 32.31 kg of head kg⁻¹ of K applied was recorded with 75% RDF + *T. harzianum* (T₃) in 2016–2017, while application of 100% RDF (T₂) registered the highest PE_K of 34.33 kg of head kg⁻¹ of K applied in 2017–2018. Results of PE_K further indicated that microbial consortium treatments gave an equivalent effect to that of single-species bio-priming treatments. Sole application of *T. harzianum* resulted in higher PE_K than other bio-primed treatments including dual and triple consortiums.

In the present study, apparent recovery efficiency (ARE_K) ranged from 59.13 to 116.49%, with a mean value of 89.34% (Table 3). The maximum ARE_K was achieved under 75% RDF + *T. harzianum* + *P. fluorescens* (T₆), which increased from 109.26% in the first year to 116.49% in the second year. Among single bio-priming agents, *P. fluorescens* showed the highest ARE_K . This microbe recorded a 13% increment in ARE_K as compared to the first year. Results followed the trend of 75% RDF + dual consortium > 75% RDF + single bio-priming agents > 75% RDF + triple consortium > 100% RDF. Application of triple consortium reduced the ARE_K by 45% (average of two years) compared to that of dual consortium (*T. harzianum* + *P. fluorescens*). Enhanced K uses efficiency in AE_K , PE_K , and ARE_K due to bacterial inoculations and reduced chemical fertilizer quantity reported by Khang-hah et al.³⁸.

Yield attributes. The observation on heading percentage and head diameter indicated that the sole application of mineral fertilizers and integrated application along with bio-agents did not bring any significant variation among the treated plants (Fig. 2). Heading percentage (72.92), as well as head diameter (13.85), was recorded to be maximum in 75% RDF + *T. harzianum* + *P. fluorescens* (T₆). The lowest heading percentage was noticed in absolute control (29.17) followed by a triple consortium (64.06). The average diameter of marketable heads was 13.35 cm. Head weight varied from 371.63 to 736.29 g (Fig. 2).

On a pooled basis, a significantly higher head weight (725.51 g) was detected with T₆ compared to the rest of the treatments. It registered 12% and 83% increments over 100% RDF (T₂) and absolute control (T₁), respectively. Bio-priming with triple consortium recorded 17% lower head weight than dual consortium (*T. harzianum* + *P. fluorescens*). Results on head weight followed the order of 75% RDF + *T. harzianum* + *P. fluorescens* (T₆) > 75% RDF + *P. fluorescens* + *B. subtilis* (T₇) > 75% RDF + *T. harzianum* + *B. subtilis* (T₈) > 100% RDF (T₂) > 75% RDF + *P. fluorescens* (T₄) > 75% RDF + *T. harzianum* (T₃) > 75% RDF + *B. subtilis* (T₅) > 75% RDF + *T. harzianum* + *P.*




Figure 2. Effect of bio-priming and fertilization on yield attributes of red cabbage. Bars (mean \pm SE; $n = 3$) followed by different alphabets significantly differ ($P \leq 0.05$) among the treatments as per DMRT. Treatments: T₁: Absolute control N:P₂O₅:K₂O @ 0:0:0 kg ha⁻¹; T₂: RDF of N:P₂O₅:K₂O @ 120:60:60 kg ha⁻¹; T₃: 75% RDF + *Trichoderma harzianum*; T₄: 75% RDF + *Pseudomonas fluorescens*; T₅: 75% RDF + *Bacillus subtilis*; T₆: 75% RDF + *T. harzianum* + *P. fluorescens*; T₇: 75% RDF + *P. fluorescens* + *B. subtilis*; T₈: 75% RDF + *T. harzianum* + *B. subtilis*; T₉: 75% RDF + *T. harzianum* + *P. fluorescens* + *B. subtilis*.

fluorescens + *B. subtilis* (T₉) > absolute control (T₁). A perusal of pooled data presented in Fig. 2 revealed that the total weight varied from 693.33 to 1030.49 g. In total weight, T₆ was 8% and 49% higher than T₂ and T₁, respectively. The total weight of plants under T₆ was at par with T₇. *P. fluorescens* combined with 75% RDF resulted in an equivalent total weight to that of 100% RDF. The direct impact of bio-priming on plant growth promotion was reported in earlier studies^{5,8}. However, these studies were conducted in pot conditions. The current study focuses on the practical utility of such technology under field conditions. Higher yield attributes in bio-primed treatments could alter cellular mechanisms in these plants and regulated nutrient supply from the soil. Integrated application of inorganic fertilizers (75% RDF) and organics (biofertilizer + vermicompost) yielded the highest marketable head percentage and a head weight of cabbage³⁹. Application of organics (poultry manure) alone showed greater head weight (17%) and head length (8%) of cabbage over NPK fertilizers⁴⁰.

Energy budgeting. Efficient use of energy inputs in a production system must lessen our dependence on non-renewable energy and maintain sustainability. The magnitude of energy input ranged from 6323 to 14,663 MJ ha⁻¹ (Table 4). Bio-priming intervention in red cabbage cultivation saved 1906 to 1965 MJ ha⁻¹ energy requirement. As expected, the energy consumed under 100% RDF (T₂) was ~15% higher than bio-priming treatments. Energy output (26,370 and 26,630 MJ ha⁻¹) and energy balance (13,643 and 13,903 MJ ha⁻¹) were considerably higher in T₆ treatment (75% RDF + *T. harzianum* + *P. fluorescens*) during both the years of study. Compared to the first year, energy balance declined (2%) in T₂ treatment. Among the treatments with amendments, T₆ showed the highest energy use efficiency (2.09). The triple consortium treatment (T₆) recorded the lowest energy balance (9799 MJ ha⁻¹) and energy use efficiency (1.77) compared to other bio-priming treatments. However, the bio-priming treatments were greater in energy balance and energy use efficiency than sole use of chemical fertilization. It shows that fertilizer inputs consume the highest energy. In cabbage production, the consumption of energy by mineral fertilizers may be as high as 77%²⁰. It is of great concern, and the non-renewable sources of plant nutrients must be substituted with renewable sources. Mihov et al.⁴¹ reported that cabbage's organic production could save 31.23% energy unit area⁻¹ than its conventional production system. Application of biofertilizers and reduced dose of fertilizers enhance energy balance and energy use efficiency of a cropping system⁴²⁻⁴⁸.

Economic analysis. Farmers will adopt any technology when it is economically feasible. Different economic indicators, such as gross return, the net return, and benefit:cost (B:C) ratio showed wide variations among the treatments (Table 5). Maximum gross return (US \$ 16,030 and 15,966 ha⁻¹) and net return (US \$ 13,877 and 13,813 ha⁻¹) were recorded with the application of T₆ treatment (75% RDF + *T. harzianum* + *P. fluorescens*) during both the years of study. The treatment also achieved the highest B:C ratio of 6.45 and 6.42. Lowest returns and B:C ratio were observed in control. Our results showed that the application of microbial consortium and 75% RDF could be profitable than 100% RDF by providing about US \$ 3222.5 higher net return on a hectare

Treatments	2016–2017				2017–2018			
	Input energy (MJ ha ⁻¹)	Output energy (MJ ha ⁻¹)	Energy balance (MJ ha ⁻¹)	Energy use efficiency	Input energy (MJ ha ⁻¹)	Output energy (MJ ha ⁻¹)	Energy balance (MJ ha ⁻¹)	Energy use efficiency
T ₁ : Absolute control N:P ₂ O ₅ :K ₂ O @ 0:0:0 kg ha ⁻¹	6323	13553 ^d	7230 ^d	2.14 ^a	6323	12187 ^c	5863 ^d	1.93 ^{ab}
T ₂ : RDF of N:P ₂ O ₅ :K ₂ O @ 120:60:60 kg ha ⁻¹	14,663	23137 ^{abc}	8473 ^{cd}	1.58 ^e	14,663	22967 ^b	8303 ^{cd}	1.57 ^c
T ₃ : 75% RDF + <i>Trichoderma harzianum</i>	12,698	22307 ^{bc}	9609 ^{bcd}	1.76 ^{cde}	12,698	23430 ^b	10732 ^{bc}	1.85 ^{ab}
T ₄ : 75% RDF + <i>Pseudomonas fluorescens</i>	12,698	23477 ^{abc}	10779 ^{abc}	1.85 ^{bcd}	12,698	23947 ^{ab}	11249 ^{abc}	1.89 ^{ab}
T ₅ : 75% RDF + <i>Bacillus subtilis</i>	12,698	22207 ^{bc}	9509 ^{bcd}	1.75 ^{cde}	12,698	22790 ^b	10092 ^{bc}	1.79 ^{bc}
T ₆ : 75% RDF + <i>T. harzianum</i> + <i>P. fluorescens</i>	12,727	26370 ^a	13643 ^a	2.07 ^{ab}	12,727	26630 ^a	13903 ^a	2.09 ^a
T ₇ : 75% RDF + <i>P. fluorescens</i> + <i>B. subtilis</i>	12,727	25303 ^{ab}	12576 ^{ab}	1.99 ^{abc}	12,727	24867 ^{ab}	12139 ^{ab}	1.95 ^{ab}
T ₈ : 75% RDF + <i>T. harzianum</i> + <i>B. subtilis</i>	12,727	23770 ^{abc}	11043 ^{abc}	1.87 ^{bcd}	12,727	24657 ^{ab}	11929 ^{ab}	1.94 ^{ab}
T ₉ : 75% RDF + <i>T. harzianum</i> + <i>P. fluorescens</i> + <i>B. subtilis</i>	12,757	21640 ^c	8883 ^{cd}	1.70 ^{de}	12,757	22557 ^b	9799 ^{bc}	1.77 ^{bc}

Table 4. Energetics of red cabbage production as influenced by bio-priming and mineral fertilization. Different letters indicate significant differences at $P \leq 0.05$ among the treatments as per DMRT.


Treatments	2016–2017			2017–2018		
	Gross return (US \$ ha ⁻¹)	Net return (US \$ ha ⁻¹)	B:C ratio	Gross return (US \$ ha ⁻¹)	Net return (US \$ ha ⁻¹)	B:C ratio
T ₁ : Absolute control N:P ₂ O ₅ :K ₂ O @ 0:0:0 kg ha ⁻¹	4098 ^c	2011 ^c	0.96 ^c	2714 ^c	627 ^c	0.30 ^c
T ₂ : RDF of N:P ₂ O ₅ :K ₂ O @ 120:60:60 kg ha ⁻¹	12743 ^{ab}	10588 ^{ab}	4.91 ^{ab}	12812 ^b	10657 ^b	4.95 ^b
T ₃ : 75% RDF + <i>Trichoderma harzianum</i>	12350 ^b	10200 ^b	4.74 ^b	12987 ^b	10836 ^b	5.04 ^b
T ₄ : 75% RDF + <i>Pseudomonas fluorescens</i>	13050 ^{ab}	10900 ^{ab}	5.07 ^{ab}	13522 ^{ab}	11372 ^{ab}	5.29 ^{ab}
T ₅ : 75% RDF + <i>Bacillus subtilis</i>	11932 ^b	9782 ^b	4.55 ^b	12667 ^b	10517 ^b	4.89 ^b
T ₆ : 75% RDF + <i>T. harzianum</i> + <i>P. fluorescens</i>	16030 ^a	13877 ^a	6.45 ^a	15966 ^a	13813 ^a	6.42 ^a
T ₇ : 75% RDF + <i>P. fluorescens</i> + <i>B. subtilis</i>	14662 ^{ab}	12509 ^{ab}	5.81 ^{ab}	14700 ^{ab}	12547 ^{ab}	5.83 ^{ab}
T ₈ : 75% RDF + <i>T. harzianum</i> + <i>B. subtilis</i>	13462 ^{ab}	11308 ^{ab}	5.25 ^{ab}	14136 ^{ab}	11983 ^{ab}	5.57 ^{ab}
T ₉ : 75% RDF + <i>T. harzianum</i> + <i>P. fluorescens</i> + <i>B. subtilis</i>	11596 ^b	9440 ^b	4.38 ^b	12271 ^b	10115 ^b	4.69 ^b

Table 5. Economics of red cabbage production as influenced by bio-priming and mineral fertilization. 1 US \$ = 66 INR; Produce cost calculated @ 0.76 US \$ kg⁻¹; Spencer (Supermarket) cost = 2.42 US \$ kg⁻¹. Different letters indicate significant differences at $P \leq 0.05$ among the treatments as per DMRT.


basis. The integrated approach's performance was better in a dual consortium in most cases, while in single-species bio-priming, application of *P. fluorescens* yields a higher B:C ratio than 100% RDF. The high profitability of a system is related to higher productivity and lowering of production cost. Thakur et al.⁴⁹ noted the highest net return and B:C ratio with the conjoint application of 75% NPK and organics (organic manures + biofertilizers) in cauliflower. Another study by Kamal et al.⁵⁰ reported a significant effect on B:C ratio of hybrid cabbage production due to the combined application of bio-agents and chemical fertilizers.

Conclusion

We have evaluated different performance indicators, each depicting new viewpoints on biotechnological intervention in integrated nutrient management. Our results generated from field experiments indicated bio-priming in combination with mineral fertilization augmented the productivity, nutrient use efficiency, and profitability of red cabbage cultivation while minimizing the energy requirements. The strategy was more effective under



(a)



(b)

Figure 3. Meteorological observations during growth period of red cabbage (a) 2016–2017 (b) 2017–2018.

dual inoculation treatments over control. Bio-priming with *T. harzianum* and *P. fluorescens* emerged out to be the most suitable treatment over control for red cabbage production, followed by bacterial co-inoculations of *P. fluorescens* and *B. subtilis*. Development of low-cost technology with a high B:C ratio has prime importance for resource-poor farmers. Our investigation also has great relevance to the United Nations Sustainable Development Goals' targets by adopting cleaner and cost-effective crop production practices to ensure food security and resource use efficiency in developing countries like India.

Materials and methods

Study site. A two-year field experiment was conducted at the Vegetable Research Farm of the Institute of Agricultural Sciences, Banaras Hindu University, Varanasi, (25°26'N, 82°99'E, and 80.7 m above mean sea level), Uttar Pradesh, India, during two consecutive winter (*rabi*) seasons of 2016–17 and 2017–18. The experimental site falls under Middle Gangetic Plains (agro-ecological region) of Eastern India. Varanasi is characterized by a semi-arid and sub-humid climate with an annual average rainfall of around 1100 mm. The cold period starts in November and stays till February, designating January as the coldest month. Meteorological observations (maximum and minimum temperature, sunshine, and evaporation) during the cropping seasons are graphically presented in Fig. 3. The mean weekly maximum and minimum temperature of the two crop seasons were recorded as 30.0 °C and 7.06 °C, respectively. Sunshine duration was higher in the second growing season. The experimental soil, Gangetic alluvial in nature, is classified in Typic Ustochrept of the order Inceptisol. Salient initial physical, chemical and biological properties (Table 6) of the soil were determined before starting field

Parameter	Value		Reference
	2016–2017	2017–2018	
Sand (%)	47.65	47.63	51
Silt (%)	30.06	30.12	
Clay (%)	22.29	22.25	
Bulk density (Mg m ⁻³)	1.36 ± 0.01	1.37 ± 0.01	52
Maximum water holding capacity (%)	42.4	42.1	53
pH (1:2)	7.58 ± 0.02	7.52 ± 0.03	54
EC (dS m ⁻¹)	0.25 ± 0.01	0.26 ± 0.01	55
Exchangeable Ca ²⁺ + Mg ²⁺ [cmol (p+) kg ⁻¹]	8.72 ± 0.29	9.16 ± 0.23	56
CEC [cmol (p+) kg ⁻¹]	18.74 ± 0.39	19.86 ± 0.53	57
Organic C (g kg ⁻¹)	4.04 ± 0.16	4.15 ± 0.20	58
Available N (kg ha ⁻¹)	203.21 ± 9.22	213.25 ± 8.87	59
Available P (kg ha ⁻¹)	20.86 ± 0.44	22.35 ± 0.31	60
Available K (kg ha ⁻¹)	217.73 ± 2.42	226.69 ± 6.40	61

Table 6. Initial properties of the soil before transplanting of crop. Values are ± standard error of mean.

experiments. The initial analysis revealed that the soil was sandy loam in texture, slightly alkaline in reaction, low in organic carbon and available N content, and medium in available P and available K content.

Seed and microbial agents. Red cabbage variety F1 Hybrid Red Ruby-2 was used as a test crop in the present study. The maturity period of the crop lies between 90 and 100 days after transplanting. Truthful label seeds with 70% germination rate and 98% physical purity were used for sowing in the nursery. The variety is suitable for growing in the Eastern Zone of India. Microbes, viz., *Trichoderma harzianum* (BHU P4; GenBank accession No. MH730446), *Pseudomonas fluorescens* (OKC; GenBank accession No. JN128891), and *Bacillus subtilis* (BHHU100; GenBank accession No. JN099686) were collected from the Department of Mycology and Plant Pathology of Institute of Agricultural Sciences, Banaras Hindu University, India. These microbial strains were compatible with each other and showed particular plant growth-promoting (PGP) traits⁶².

Bio-priming. *Preparation of inoculum.* Bacterial (*B. subtilis* and *P. fluorescens*) culture was inoculated in 250 mL flasks containing 100 mL nutrient broth and kept in a shaking incubator (150 rpm; 28 ± 2 °C) for 48 h. Bacterial pellets were obtained by centrifugation (7000 rpm; 4 °C) for 10 min. Discarding the supernatant, the cell pellets were soon washed with sterile distilled water. An adjustment of final cell density to 4 × 10⁸ CFU mL⁻¹ was done using optical density (< 1) at 600 nm⁶³. In *T. harzianum*, spore suspension was prepared from 1 week of culture grown (28 ± 2 °C) on potato dextrose agar by harvesting the spores in sterilized 0.85% sodium chloride (NaCl)⁵. The spore concentration was adjusted to 2 × 10⁷ CFU mL⁻¹ by measuring the optical density in a spectrophotometer. Cell suspensions and/or spore suspension were mixed in equal ratios for the dual consortium (1:1) and triple consortium (1:1:1).

Seedling bio-priming. Red cabbage seeds were washed with tap water and surface sterilized with 0.1% mercuric chloride (HgCl₂) solution for 2 min. After sterilization, the seeds were immediately washed with autoclaved distilled water three times. Next, the seeds were soaked in sterile distilled water for 2 h. After the hydration treatment, the seeds were sown in the nursery. Seedlings were picked when there were 5–6 leaves. Soil attached with the roots was washed carefully, followed by root dipping in liquid culture containing 2% carboxymethyl cellulose (CMC) as an adhesive agent. The bio-priming process was followed for 5 h under incubated conditions (28 ± 2 °C; > 90% relative humidity)^{5,36,62}.

Experimental design and crop management. The field setup was laid out in a randomized block design with nine treatments and three replications. Red cabbage seeds were sown in raised bed nursery about 1 month before transplanting. The field was twice ploughed by a tractor and planked 15 days before the implementation of the experiment. Crop residues, stones, pebbles, or weeds were removed manually from the field. Five weeks of healthy and uniform-sized red cabbage seedlings were transplanted in 4 × 2 m² plots with a spacing of 50 cm × 50 cm on 5th December 2016 and 2017. Four rows in a plot accommodated 32 plants. The crop was irrigated for 2 weeks with a watering can just after transplanting. After that, it received four irrigations at 7–10 days intervals as per the requirement. A full dose of diammonium phosphate (DAP) and muriate of potash (MOP) was applied as basal at the time of final land preparation. The urea dose was given in three splits, including basal application and top dressing, at 30 and 45 days (head initiation) after transplanting plant seedlings. Other intercultural operations like gap filling and weeding (3 times) were also carried. In general, the pest and disease incidence was not observed during both crop seasons. The treatments comprised nine combinations of mineral fertilizers and bio-agents, including one absolute control outlined in Table 7. The recommended dose of fertilizer (RDF) was applied @ 120:60:60 kg ha⁻¹ (N:P₂O₅:K₂O) through urea, DAP, and MOP, respectively. The fertilizer dose was reduced to 25% when the priming agents were used (Table 7).

Treatments	Notations used	Mineral fertilization (kg ha ⁻¹)		
		N	P ₂ O ₅	K ₂ O
Absolute control N:P ₂ O ₅ :K ₂ O @ 0:0:0 kg ha ⁻¹	T ₁	0	0	0
RDF of N:P ₂ O ₅ :K ₂ O @ 120:60:60 kg ha ⁻¹	T ₂	120	60	60
75% RDF + <i>Trichoderma harzianum</i>	T ₃	90	45	45
75% RDF + <i>Pseudomonas fluorescens</i>	T ₄	90	45	45
75% RDF + <i>Bacillus subtilis</i>	T ₅	90	45	45
75% RDF + <i>T. harzianum</i> + <i>P. fluorescens</i>	T ₆	90	45	45
75% RDF + <i>P. fluorescens</i> + <i>B. subtilis</i>	T ₇	90	45	45
75% RDF + <i>T. harzianum</i> + <i>B. subtilis</i>	T ₈	90	45	45
75% RDF + <i>T. harzianum</i> + <i>P. fluorescens</i> + <i>B. subtilis</i>	T ₉	90	45	45

Table 7. Details of treatment combinations applied during the study.

Biometric observations. Data related to yield attributing parameters such as heading percentage, head diameter (cm), head weight (g), and total weight (g) were collected at harvest.

Computation of nutrient uptake and nutrient use efficiency. Nutrient uptake and nutrient use efficiency were calculated using the following equations.

Nutrient uptake.

$$\text{Nutrient uptake (kg ha}^{-1}\text{)} = \frac{\text{Nutrient content (\%)} \times \text{Yield (kg ha}^{-1}\text{)}}{100}$$

Nitrogen content was determined by the micro Kjeldahl method after pre-digestion with concentrated sulphuric acid (H₂SO₄) followed by catalyst mixture⁶⁴. Phosphorus content was determined by vanadomolybdophosphoric yellow color method after pre-digestion with diacid (HNO₃:HClO₄)⁶⁵. The acid-digest prepared for P was used to assess potassium (K) in a flame photometer⁶⁶.

Agronomic efficiency.

$$\text{Agronomic efficiency (kg/ha)} = \frac{Y - Y_0}{\text{Quantity (kg/ha) of nutrients applied}} \times 100$$

where Y = head yield (kg ha⁻¹) with applied nutrient, Y₀ = head yield (kg ha⁻¹) with no applied nutrient.

Physiological efficiency.

$$\text{Physiological efficiency (kg kg}^{-1}\text{)} = \frac{Y - Y_0}{U - U_0}$$

where, Y = head yield (kg ha⁻¹) with applied nutrient, Y₀ = head yield (kg ha⁻¹) with no applied nutrient, U = total nutrient uptake (kg ha⁻¹) with applied nutrient; U₀ = total nutrient uptake (kg ha⁻¹) with no applied nutrient.

Apparent recovery efficiency.

$$\text{Apparent recovery efficiency (\%)} = \frac{U - U_0}{\text{Quantity (kg ha}^{-1}\text{) of nutrient applied}} \times 100$$

where, U = total nutrient uptake (kg ha⁻¹) with applied nutrient; U₀ = total nutrient uptake (kg ha⁻¹) with no applied nutrient.

Partial factor productivity.

$$\text{Partial factor productivity (kg kg}^{-1}\text{)} = \frac{\text{Head yield (kg ha}^{-1}\text{)}}{\text{Quantity (kg ha}^{-1}\text{) of nutrient applied}}$$

Energy analysis. The energy input–output relationship was determined based on the energy equivalent of inputs and output (Table 8). Solar energy was not included in the calculation. Energy analysis involved the following equations.

Particulars	Units	Energy equivalent (MJ unit ⁻¹) <small>20,21,41</small>	Unit cost (US \$ unit ⁻¹)
Human labor	hr	1.96	1.26
Diesel	L	56.31	0.98
Farm machinery	hr	62.70	7.57
Electricity	kWh	11.91	
Mineral fertilizer			
N	kg	60.60	0.227
P ₂ O ₅	kg	11.10	0.378
K ₂ O	kg	6.70	0.303
Biofertilizer	L	2.98	7.57
Irrigation water	m ³	1.02	0.0076
Seed	kg	0.8	454
Produce	kg	0.8	0.76

Table 8. Energy equivalents and cost of inputs and outputs in red cabbage production. 1 US \$ = 66 INR.

$$\text{Energy use efficiency} = \frac{\text{Energy output (MJ ha}^{-1}\text{)}}{\text{Energy input (MJ ha}^{-1}\text{)}}$$

$$\text{Energy productivity} = \frac{\text{Head yield (kg ha}^{-1}\text{)}}{\text{Energy input (MJ ha}^{-1}\text{)}}$$

$$\text{Energy balance} = \text{Energy output (MJ ha}^{-1}\text{)} - \text{Energy input (MJ a}^{-1}\text{)}$$

Economic analysis. Expenditures were calculated considering the unit cost of variable inputs and output (produce) based on prevailing market prices (Table 8). The cost of cultivation (total expenditure) was subtracted from the gross return for obtaining the net return. Benefit: cost (B:C) ratio was worked out using the following formula.

$$\text{B : Cratio} = \frac{\text{Net return (US \$ ha}^{-1}\text{)}}{\text{Total expenditure (US \$ ha}^{-1}\text{)}}$$

Statistical analysis. Data collected during the study were subjected to one-way analysis of variance (ANOVA). Values are presented as mean \pm standard error (SE). Significance of difference between treatment means was performed using Duncan's multiple range test (DMRT) at $P \leq 0.05$ significance level. Statistical Package for Social Science (SPSS, version 20) software was used for these analyses^{67,68}.

International, national and/or institutional guidelines. Authors reporting experiments confirmed that the use of plants in the present study complies with international, national and/or institutional guidelines.

Received: 11 March 2021; Accepted: 15 July 2021

Published online: 03 August 2021

References.

- Ramakrishna-Parama, V. R. Need for sustained nutrient cycling and management—issues and responses. *J. Indian Soc. Soil Sci.* **62**, S86–S94 (2014).
- Priyadarshini, P. & Abhilash, P. C. Policy recommendations for enabling transition towards sustainable agriculture in India. *Land Use Policy* **96**, 104718 (2020).
- Sarkar, D. & Rakshit, A. Safeguarding the fragile rice–wheat ecosystem of the Indo-Gangetic Plains through bio-priming and bioaugmentation interventions. *FEMS Microbiol. Ecol.* **96**, faa221 (2020).
- Meena, S. K., Rakshit, A. & Meena, V. S. Effect of seed bio-priming and N doses under varied soil type on nitrogen use efficiency (NUE) of wheat (*Triticum aestivum* L.) under greenhouse conditions. *Biocatal. Agric. Biotechnol.* **6**, 68–75 (2016).
- Singh, V., Upadhyay, R. S., Sarma, B. K. & Singh, H. B. *Trichoderma asperellum* spore dose depended modulation of plant growth in vegetable crops. *Microbiol. Res.* **193**, 74–86 (2016).
- Chatterjee, N. *et al.* On-farm seed priming interventions in agronomic crops. *Acta Agric. Slov.* **111**, 715–735 (2018).
- Yadav, S. K. *et al.* Co-inoculated biopriming with *Trichoderma*, *Pseudomonas* and *Rhizobium* improves crop growth in *Cicer arietinum* and *Phaseolus vulgaris*. *Int. J. Agric. Environ. Biotechnol.* **6**, 255–259 (2013).
- Meena, S. K., Rakshit, A., Singh, H. B. & Meena, V. S. Effect of nitrogen levels and seed bio-priming on root infection, growth and yield attributes of wheat in varied soil type. *Biocatal. Agric. Biotechnol.* **12**, 172–178 (2017).

9. Bottner, P., Pansu, M. & Sallih, Z. Modelling the effect of active roots on soil organic matter turnover. *Plant Soil* **216**, 15–25 (1999).
10. Buchenauer, H. Biological control of soil-borne diseases by rhizobacteria/Biologische Bekämpfung von bodenbürtigen Krankheiten durch Rhizobakterien. *Zeitschrift für Pflanzenkrankheiten und Pflanzenschutz/Journal Plant Dis. Prot.* **105**, 329–348 (1998).
11. Glick, B. R. Plant growth-promoting bacteria: mechanisms and applications. *Scientifica (Cairo)*. **2012**, 963401 (2012).
12. Ahemad, M. & Kibret, M. Mechanisms and applications of plant growth promoting rhizobacteria: current perspective. *J. King Saud Univ. Sci.* **26**, 1–20 (2014).
13. de Souza, R., Ambrosini, A. & Passaglia, L. M. P. Plant growth-promoting bacteria as inoculants in agricultural soils. *Genet. Mol. Biol.* **38**, 401–419 (2015).
14. Sarkar, D., Pal, S., Singh, H. B., Yadav, R. S. & Rakshit, A. Harnessing Bio-priming for Integrated Resource Management under Changing Climate. In *Advances in PGPR Research vol 349* (eds Singh, H. B. et al.) (CABI International, 2017).
15. Rakshit, A. Impact assessment of bio priming mediated nutrient use efficiency for climate resilient agriculture. In *Climate Change and Agriculture in India: Impact and Adaptation 57–68* (Springer, 2019).
16. Paungfoo-Lonhienne, C., Redding, M., Pratt, C. & Wang, W. Plant growth promoting rhizobacteria increase the efficiency of fertilisers while reducing nitrogen loss. *J. Environ. Manag.* **233**, 337–341 (2019).
17. Pereira, S. I. A., Abreu, D., Moreira, H., Vega, A. & Castro, P. M. L. Plant growth-promoting rhizobacteria (PGPR) improve the growth and nutrient use efficiency in maize (*Zea mays* L.) under water deficit conditions. *Heliyon* **6**, e05106 (2020).
18. Yadav, G. S. *et al.* Energy budgeting for designing sustainable and environmentally clean/safer cropping systems for rainfed rice fallow lands in India. *J. Clean. Prod.* **158**, 29–37 (2017).
19. Singh, K. P., Prakash, V., Srinivas, K. & Srivastva, A. K. Effect of tillage management on energy-use efficiency and economics of soybean (*Glycine max*) based cropping systems under the rainfed conditions in North-West Himalayan Region. *Soil Tillage Res.* **100**, 78–82 (2008).
20. Tuti, M. D. *et al.* Energy budgeting of colocasia-based cropping systems in the Indian sub-Himalayas. *Energy* **45**, 986–993 (2012).
21. Mihov, M. & Tringovska, I. Energy efficiency improvement of greenhouse tomato production by applying new biofertilizers. *Bulg. J. Agric. Sci.* **16**, 454–458 (2010).
22. Pal, S. & Singh, H. B. Energy inputs and yield relationship in greenhouse okra production by bio-priming. *Int. J. Agric. Environ. Biotechnol.* **11**, 741–746 (2018).
23. Mizgier, P. *et al.* Characterization of phenolic compounds and antioxidant and anti-inflammatory properties of red cabbage and purple carrot extracts. *J. Funct. Foods* **21**, 133–146 (2016).
24. Sarkar, D. & Rakshit, A. Red cabbage as potential functional food in the present perspective. *Int. J. Bioresour. Sci.* **4**, 7 (2017).
25. Narayanamma, M., Chiranjeevi, C. H., Reddy, I. P. & Ahmed, S. R. Integrated nutrient management in cauliflower (*Brassica oleracea* var botrytis L.). *Veg Sci* **32**, 62–64 (2005).
26. Chatterjee, R., Choudhuri, P. & Thirumdasu, R. K. Uptake and availability of major nutrients in cabbage crop amended with organic and inorganic nutrient sources under Eastern Himalayan Region. *J. Basic Appl. Res. Int.* **2**, 100–105 (2016).
27. Bhardwaj, S., Kaushal, R., Kaushal, M. & Bhardwaj, K. K. Integrated nutrient management for improved cauliflower yield and soil health. *Int. J. Veg. Sci.* **24**, 29–42 (2018).
28. Danish, S., Zafar-Ul-Hye, M., Hussain, S., Riaz, M. & Qayyum, M. F. Mitigation of drought stress in maize through inoculation with drought tolerant ACC deaminase containing PGPR under axenic conditions. *Pakistan J. Bot.* **52**, 49–60 (2020).
29. Danish, S., Zafar-ul-Hye, M., Mohsin, F. & Hussain, M. ACC-deaminase producing plant growth promoting rhizobacteria and biochar mitigate adverse effects of drought stress on maize growth. *PLoS ONE* **15**, e0230615 (2020).
30. Zafar-ul-Hye, M., Danish, S., Abbas, M., Ahmad, M. & Munir, T. M. ACC deaminase producing PGPR *Bacillus amyloliquefaciens* and *Agrobacterium fabrum* along with biochar improve wheat productivity under drought stress. *Agron.* **9**, 343 (2019).
31. Danish, S. *et al.* Drought stress alleviation by ACC deaminase producing *Achromobacter xylosoxidans* and *Enterobacter cloacae*, with and without timber waste biochar in maize. *Sustainability* **12**, 6286 (2020).
32. Kaur, G. & Reddy, M. S. Role of phosphate-solubilizing bacteria in improving the soil fertility and crop productivity in organic farming. *Arch. Agron. Soil Sci.* **60**, 549–564 (2014).
33. Basak, N. *et al.* Impact of long-term application of organics, biological, and inorganic fertilizers on microbial activities in rice-based cropping system. *Commun. Soil Sci. Plant Anal.* **48**, 2390–2401 (2017).
34. Ranjit, C., Bandhopadhyay, S. & Jana, J. C. Organic amendments influencing growth, head yield and nitrogen use efficiency in cabbage (*Brassica oleracea* var Capitata L.). *Am. Int. J. Res. Formal Appl. Nat. Sci.* **5**, 90–95 (2014).
35. Visconti, D. *et al.* Can Trichoderma-based biostimulants optimize N Use efficiency and stimulate growth of leafy vegetables in greenhouse intensive cropping systems? *Agronomy* **10**, 121 (2020).
36. Yaseen, M. & Malhi, S. S. Variation in yield, phosphorus uptake, and physiological efficiency of wheat genotypes at adequate and stress phosphorus levels in soil. *Commun. Soil Sci. Plant Anal.* **40**, 3104–3120 (2009).
37. Emami, S. *et al.* Consortium of endophyte and rhizosphere phosphate solubilizing bacteria improves phosphorous use efficiency in wheat cultivars in phosphorus deficient soils. *Rhizosphere* **14**, e100196 (2020).
38. Khanghahi, M. Y., Pirdashti, H., Rahimian, H., Nematzadeh, G. & Sepanlou, M. G. Potassium solubilising bacteria (KSB) isolated from rice paddy soil: from isolation, identification to K use efficiency. *Symbiosis* **76**, 13–23 (2018).
39. Chatterjee, R. Physiological attributes of cabbage (*Brassica oleracea*) as influenced by different sources of nutrients under eastern Himalayan region. *Res. J. Agric. Sci.* **1**, 318–321 (2010).
40. Moyin-Jesu, E.I. Use of different organic fertilizers on soil fertility improvement, growth and head yield parameters of cabbage (*Brassica oleracea* L.). *Intnl. J. Recycl. Organic Waste Agri.* **4**, 291–298.
41. Mihov, M. *et al.* Energy assessment of conventional and organic production of head cabbage. *Bulg. J. Agric. Sci.* **18**, 320–324 (2012).
42. Munda, S. *et al.* Inorganic phosphorus along with biofertilizers improves profitability and sustainability in soybean (*Glycine max*)-potato (*Solanum tuberosum*) cropping system. *J. Saudi Soc. Agric. Sci.* **17**, 107–113 (2018).
43. Tahir, F. A., Ahamad, N., Rasheed, M. K. & Danish, S. Effect of various application rate of zinc fertilizer with and without fruit waste biochar on the growth and Zn uptake in maize. *Int. J. Biosci.* **13**, 159–166 (2018).
44. Ahmed, N. *et al.* Rhizobacteria and silicon synergy modulates the growth, nutrition and yield of mungbean under saline soil. *Pakistan J. Bot.* **52**, 9–15 (2020).
45. Adnan, M. *et al.* Coupling phosphate-solubilizing bacteria with phosphorus supplements improve maize phosphorus acquisition and growth under lime induced salinity stress. *Plants* **9**, 900 (2020).
46. Rafullah, *et al.* Phosphorus nutrient management through synchronization of application methods and rates in wheat and maize crops. *Plants* **9**, 1389 (2020).
47. Zafar-ul-Hye, M., Hussain, N. M., Danish, S., Aslam, U. & Zahir, Z. A. Multi-Strain bacterial inoculation of *Enterobacter cloacae*, *Serratia ficaria* and *Burkholderia phytofirmans* with fertilizers for enhancing resistance in wheat against salinity stress. *Pakistan J. Bot.* **51**, 1839–1846 (2019).
48. Zafar-ul-Hye, M., Zahra, M. B., Danish, S. & Abbas, M. Multi-strain inoculation with PGPR producing ACC deaminase is more effective than single-strain inoculation to improve wheat (*Triticum aestivum*) growth and yield. *Phyton-Int. J. Exp. Bot.* **89**, 405–413 (2020).
49. Thakur, J., Kumar, P. & Mohit. Studies on conjoint application of nutrient sources and PGPR on growth, yield, quality, and economics of cauliflower (*Brassica oleracea* var. botrytis L.). *J. Plant Nutr.* **41**, 1862–1867 (2018).

50. Kamal, K., Devi, S. & Prasad V.M. Effect of microbial inoculants and chemical fertilizers on yield and economics of hybrid cabbage (*Brassica oleracea* var. capitata). *Asian J. Hortic.* **11**, 338–343 (2016).
51. Gee, G. W. & Bauder, J. W. Particle-size analysis. In *Methods of soil analysis. Part 1. Physical and mineralogical methods* 383–411 (1986). <https://doi.org/10.2136/sssabookser5.1.2ed.c15>.
52. Blake, G. R. Bulk Density. in *Methods of Soil Analysis: Part 1 Physical and Mineralogical Properties, Including Statistics of Measurement and Sampling, 9.1* (eds. Black, C. A. & Evans, D. D.) 374–390 (American Society of Agronomy, Crop Science Society of America, and Soil Science Society of America, 1965).
53. Klute, A. Water Capacity. In *Methods of Soil Analysis: Part 1 Physical and Mineralogical Properties, Including Statistics of Measurement and Sampling, 9.1* (eds. Black, C. A. & Evans, D. D.) 273–278 (American Society of Agronomy, Crop Science Society of America, and Soil Science Society of America, 1965).
54. Page, A. L., Miller, R. H. & Keeny, D. R. Soil pH and lime requirement. In *Methods of Soil Analysis* 199–208 (American Society of Agronomy, 1982).
55. Rhoades, J. D. Salinity: Electrical Conductivity and Total Dissolved Solids. In *Methods of Soil Analysis, Part 3, Chemical Methods* (eds. D.L. Sparks et al.) vol. 5 417–435 (Soil Science Society of America, 1996).
56. Hesse, P. R. *A textbook of soil Chemical Analysis* (Cambridge University Press, New York, 1971).
57. Chapman, H. D. Cation-exchange capacity. In *Methods of Soil Analysis: Part 2 Chemical and Microbiological Properties, 92.* (ed. Norman, A. G.) 891–901 (Wiley, 1965). <https://doi.org/10.2134/agronmonogr9.2.c6>.
58. Walkley, A. & Black, I. A. An examination of the degtjareff method for determining soil organic matter, and a proposed modification of the chromic acid titration method. *Soil Sci.* **37**, 29–38 (1934).
59. Subbiah, B. V. & Asija, G. L. A rapid procedure for the determination of available nitrogen in soils. *Curr. Sci.* **25**, 259–260.
60. Olsen, S., Cole, C., Watanabe, F. & Dean, L. Estimation of available phosphorus in soils by extraction with sodium bicarbonate. No. 939. (United States Department of Agriculture, Washington, D.C. 1954).
61. Pratt, P. F. Potassium. In *Methods of Soil Analysis: Part 2 Chemical and Microbiological Properties, 9.2* (ed. Norman, A. G.) 1022–1030 (Wiley, 1965). <https://doi.org/10.2134/agronmonogr9.2.c20>.
62. Sarkar, D. & Rakshit, A. Bio-priming in combination with mineral fertilizer improves nutritional quality and yield of red cabbage under Middle Gangetic Plains India. *Sci. Hortic. (Amsterdam)* **283**, 110 (2021).
63. Jain, A., Singh, A., Singh, S. & Singh, H. B. Phenols enhancement effect of microbial consortium in pea plants restrains *Sclerotinia sclerotiorum*. *Biol. Control* **89**, 23–32 (2015).
64. Donald, A. H. & Miller, R. O. Determination of total nitrogen in plant tissue. In *Handbook of Reference Methods for Plant Analysis* (ed. Kalra, Y.) 75–83 (CRC Press, 1998).
65. Miller, O. Nitric-perchloric acid wet digestion in an open vessel. In *Reference Methods for Plant Analysis* (ed. Kalra, Y.) 57–62 (CRC Press, New York, 1998).
66. Donald, A. H. & Hanson, D. Determination of potassium and sodium by flame emission spectrophotometry. In *Handbook of Reference Methods for Plant Analysis* (ed. Kalra, Y.) 153–155 (CRC Press, 1998).
67. Steel, R. G., Torrie, J. H. & Dickey, D. A. *Principles and Procedures of Statistics: A Biometrical Approach* (McGraw Hill Book International Co., 1997).
68. S.P.S.S. *SPSS Statistics for Windows, Version 20.0.* (SPSS Inc).

Acknowledgements

We acknowledge the support received from the Head, Department of Soil Science and Agricultural Chemistry, BHU, in fulfilling the research. Deepranjan Sarkar is grateful to BHU for providing the Ph.D. Research Fellowship.

Author contributions

Conceptualization, A.R.; D.S.; methodology, D.S.; A.S.; O.S.D.; S.S.; software, D.S.; validation, A.R.; M.P.; R.Z.S.; data curation, S.; A.S.; writing—original draft preparation, D.S.; A.S.; O.S.D.; S.S.; S.; M.P.; A.R., S.D.; S.F.; R.D.; writing—review and editing, D.S., S.D.; S.F.; R.D.; R.Z.S.; A.G.; M.J.A.; visualization, D.S.; S.D.; S.F.; R.D.; Supervision, A.R.

Competing interests

The authors declare no competing interests.

Additional information

Supplementary Information The online version contains supplementary material available at <https://doi.org/10.1038/s41598-021-95092-6>.

Correspondence and requests for materials should be addressed to A.R. or S.D.

Reprints and permissions information is available at www.nature.com/reprints.

Publisher's note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.



Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

© The Author(s) 2021